

SOME ASPECTS OF LARGE TIME BEHAVIOR OF THE HEAT KERNEL: AN OVERVIEW WITH PERSPECTIVES

YEHUDA PINCHOVER

Dedicated to Professor Michael Demuth on the occasion of his 65th birthday

ABSTRACT. We discuss a variety of developments in the study of large time behavior of the positive minimal heat kernel of a time independent (not necessarily symmetric) second-order parabolic operator defined on a domain $M \subset \mathbb{R}^d$, or more generally, on a noncompact Riemannian manifold M . Our attention is mainly focused on *general* results in general settings.

CONTENTS

1. Introduction	1
2. Preliminaries	6
3. Capacitory potential and heat content	13
4. Varadhan's lemma	15
5. Existence of $\lim_{t \rightarrow \infty} e^{\lambda_0 t} k_P^M(x, y, t)$	17
6. Applications of Theorem 1.3	21
7. Davies' conjecture concerning strong ratio limit	26
8. Comparing decay of critical and subcritical heat kernels	34
9. On the equivalence of heat kernels	40
References	43

1. INTRODUCTION

The large time behavior of the heat kernel of a second-order parabolic operator has been extensively studied over the recent decades (see for example the following monographs and survey articles [6, 11, 17, 18,

Date: August 31, 2012.

2000 *Mathematics Subject Classification.* Primary 35K08; Secondary 35B09, 47D07, 47D08.

Key words and phrases. Heat kernel, Green function, parabolic Martin boundary, positive solutions, ratio limit.

20, 25, 29, 37, 54, 55, 57, 59, 62, 63, 64], and references therein). The purpose of the present paper is to review a variety of developments in this area, and to point out a number of their consequences. Our attention is mainly focused on *general* results in general settings. Still, the selection of topics in this survey is incomplete, and is according to the author's working experience and taste. The reference list is far from being complete and serves only this exposé.

Let P be a general linear, second-order, elliptic operator defined on a domain $M \subset \mathbb{R}^d$ or, more generally, on a noncompact, connected, Riemannian manifold M of dimension $d \geq 1$. Denote the cone of all positive solutions of the equation $Pu = 0$ in M by $\mathcal{C}_P(M)$. The *generalized principal eigenvalue* is defined by

$$\lambda_0 = \lambda_0(P, M) := \sup\{\lambda \in \mathbb{R} \mid \mathcal{C}_{P-\lambda}(M) \neq \emptyset\}.$$

Throughout this paper we always assume that $\lambda_0 > -\infty$.

Suppose that $\lambda_0 \geq 0$, and consider the (time-independent) parabolic operator

$$(1.1) \quad Lu := \partial_t u + P(x, \partial_x)u \quad (x, t) \in M \times (0, \infty).$$

We denote by $\mathcal{H}_P(M \times (a, b))$ the cone of all nonnegative solutions of the parabolic equation

$$(1.2) \quad Lu = 0 \quad \text{in } M \times (a, b).$$

Let $k_P^M(x, y, t)$ be the *positive minimal heat kernel* of the parabolic operator L on the manifold M . By definition, for a fixed $y \in M$, the function $(x, t) \mapsto k_P^M(x, y, t)$ is the minimal positive solution of the equation

$$(1.3) \quad Lu = 0 \quad \text{in } M \times (0, \infty),$$

subject to the initial data δ_y , the Dirac distribution at $y \in M$. It can be easily checked that for $\lambda \leq \lambda_0$, the heat kernel $k_{P-\lambda}^M$ of the operator $P - \lambda$ on M satisfies the identity

$$(1.4) \quad k_{P-\lambda}^M(x, y, t) = e^{\lambda t} k_P^M(x, y, t).$$

So, it is enough to study the large time behavior of $k_{P-\lambda_0}^M$, and therefore, in most cases we assume that $\lambda_0(P, M) = 0$. Note that the heat kernel of the operator P^* , the formal adjoint of the operator P on M , satisfies the relation

$$k_{P^*}^M(x, y, t) = k_P^M(y, x, t).$$

Here we should mention that many authors derived upper and lower Gaussian bounds for heat kernels of elliptic operators on $M := \mathbb{R}^d$, or more generally, on noncompact Riemannian manifolds M . As a

prototype result, let us recall the following classical result of Aronson [6]:

Example 1.1. Let P be a second-order uniformly elliptic operator in divergence form on \mathbb{R}^d with real coefficients satisfying some general boundedness assumptions. Then the following Gaussian estimates hold:

$$\begin{aligned} C_1(4\pi t)^{-d/2} \exp\left(-C_2 \frac{|x-y|^2}{t} - \omega_1 t\right) &\leq k_P^{\mathbb{R}^d}(x, y, t) \\ &\leq C_3(4\pi t)^{-d/2} \exp\left(-C_4 \frac{|x-y|^2}{t} + \omega_2 t\right) \quad \forall (x, y, t) \in \mathbb{R}^d \times \mathbb{R}^d \times \mathbb{R}_+. \end{aligned}$$

However, since Gaussian estimates of the above type in general are not tight as $t \rightarrow \infty$, such bounds do not provide us with the exact large time behavior of the heat kernel, let alone strong ratio limits of two heat kernels.

In spite of this, and as a first and rough result concerning the large time behavior of the heat kernel, we have the following explicit and useful formula

$$(1.5) \quad \lim_{t \rightarrow \infty} \frac{\log k_P^M(x, y, t)}{t} = -\lambda_0.$$

We note that (1.5) holds in the general case, and characterizes the generalized principal eigenvalue λ_0 in terms of the large time behavior of $\log k_P^M(x, y, t)$. The above formula is well known in the symmetric case, see for example [31] and [27, Theorem 10.24]. For the proof in the general case, see Corollary 6.1.

To get a more precise result one should introduce the notion of criticality. We say that the operator P is *subcritical* (respectively, *critical*) in M if for some $x \neq y$, and therefore for any $x \neq y, x, y \in M$, we have

$$(1.6) \quad \int_0^\infty k_P^M(x, y, \tau) d\tau < \infty \quad \left(\text{respectively, } \int_0^\infty k_P^M(x, y, \tau) d\tau = \infty \right).$$

It follows from the above definition that, roughly speaking, the heat kernel of a subcritical operator in M “decays” faster as $t \rightarrow \infty$ than the heat kernel of a critical operator in M . This rule of thumb will be discussed in Section 8.

If P is subcritical in M , then the function

$$(1.7) \quad G_P^M(x, y) := \int_0^\infty k_P^M(x, y, \tau) d\tau$$

is called the *positive minimal Green function* of the operator P in M .

It follows from (1.5) that for $\lambda < \lambda_0$, the operator $P - \lambda$ is subcritical in M . Clearly, P is subcritical (respectively, critical) in M , if and only if P^* , is subcritical (respectively, critical) in M . Furthermore, it is well known that if P is critical in M , then $\mathcal{C}_P(M)$ is a one-dimensional cone, and any positive supersolution of the equation $Pu = 0$ in M is in fact a solution. In this case, the unique positive solution $\varphi \in \mathcal{C}_P(M)$ is called *Agmon ground state* (or in short ground state) of the operator P in M [1, 48, 54]. We denote the ground state of P^* by φ^* .

The following example demonstrates two prototype behaviors as $t \rightarrow \infty$ of heat kernels corresponding to two particular classical cases.

Example 1.2. 1. Let $P = -\Delta$, $M = \mathbb{R}^d$, where $d \geq 1$. It is well known that $\lambda_0(-\Delta, \mathbb{R}^d) = 0$, and that the heat kernel is given by the Gaussian kernel. Hence,

$$\begin{aligned} e^{\lambda_0 t} k_{-\Delta}^{\mathbb{R}^d}(x, y, t) &= k_{-\Delta}^{\mathbb{R}^d}(x, y, t) \\ &= \frac{1}{(4\pi t)^{d/2}} \exp\left(\frac{-|x - y|^2}{4t}\right) \underset{t \rightarrow \infty}{\sim} t^{-d/2} \underset{t \rightarrow \infty}{\rightarrow} 0. \end{aligned}$$

So, the rate of the decay depends on the dimension, and clearly, $-\Delta$ is critical in \mathbb{R}^d if and only if $d = 1, 2$. Nevertheless, the above limit is 0 in any dimension.

2. Suppose that P is a symmetric nonnegative elliptic operator with real and smooth coefficients which is defined on a smooth bounded domain $M \Subset \mathbb{R}^d$, and let $\{\varphi_n\}_{n=0}^\infty$ be the complete orthonormal sequence of the (Dirichlet) eigenfunctions of P with the corresponding nondecreasing sequence of eigenvalues $\{\lambda_n\}_{n=0}^\infty$. Then the heat kernel has the eigenfunction expansion

$$(1.8) \quad k_P^M(x, y, t) = \sum_{n=0}^{\infty} e^{-\lambda_n t} \varphi_n(x) \varphi_n(y).$$

Hence,

$$e^{\lambda_0 t} k_P^M(x, y, t) = \sum_{n=0}^{\infty} e^{(\lambda_0 - \lambda_n)t} \varphi_n(x) \varphi_n(y) \underset{t \rightarrow \infty}{\rightarrow} \varphi_0(x) \varphi_0(y) > 0.$$

Therefore, the operator $P - \lambda_0$ is critical in M , $t^{-1} \log k_{P-\lambda_0}^M(x, y, t) \underset{t \rightarrow \infty}{\rightarrow} 0$, but $k_{P-\lambda_0}^M$ does not decay as $t \rightarrow \infty$.

3. Several explicit formulas of heat kernels of certain classical operators are included in [10, 20, 27], each of which either tends to zero or converges to a positive function as $t \rightarrow \infty$.

The characterization of λ_0 in terms of the large time behavior of the heat kernel, given by (1.5), provides us with the asymptotic behavior of $\log k_P^M$ as $t \rightarrow \infty$ but not of k_P^M itself. Moreover, (1.5) does not distinguish between critical and subcritical operators. In the first part of the present article we provide a complete proof that

$$\lim_{t \rightarrow \infty} e^{\lambda_0 t} k_P^M(x, y, t)$$

always exists. This basic result has been proved in two parts in [48] and [51], and here, for the first time, we give a comprehensive and a bit simplified proof (see also [5, 12, 28, 29, 32, 54, 60, 62, 66] and references therein for previous and related results). We have:

Theorem 1.3 ([48, 51]). *Let P be an elliptic operator defined on M , and assume that $\lambda_0(P, M) \geq 0$.*

(i): The subcritical case: *If $P - \lambda_0$ is subcritical in M , then*

$$\lim_{t \rightarrow \infty} e^{\lambda_0 t} k_P^M(x, y, t) = 0.$$

(ii): The positive-critical case: *If $P - \lambda_0$ critical is in M , and the ground states φ and φ^* of $P - \lambda_0$ and $P^* - \lambda_0$, respectively, satisfy $\varphi^* \varphi \in L^1(M)$, then*

$$\lim_{t \rightarrow \infty} e^{\lambda_0 t} k_P^M(x, y, t) = \frac{\varphi(x) \varphi^*(y)}{\int_M \varphi^*(z) \varphi(z) \, dz}.$$

(iii): The null-critical case: *If $P - \lambda_0$ is critical in M , and the ground states φ and φ^* of $P - \lambda_0$ and $P^* - \lambda_0$, respectively, satisfy $\varphi^* \varphi \notin L^1(M)$, then*

$$\lim_{t \rightarrow \infty} e^{\lambda_0 t} k_P^M(x, y, t) = 0.$$

Moreover, for $\lambda < \lambda_0$, let $G_{P-\lambda}^M(x, y)$ be the minimal positive Green function of the elliptic operator $P - \lambda$ on M . Then the following Abelian-Tauberian relation holds

$$(1.9) \quad \lim_{t \rightarrow \infty} e^{\lambda_0 t} k_P^M(x, y, t) = \lim_{\lambda \nearrow \lambda_0} (\lambda_0 - \lambda) G_{P-\lambda}^M(x, y).$$

The outline of the present paper is as follows. In Section 2 we provide a short review of the theory of positive solutions and the basic properties of the heat kernel. The proof of Theorem 1.3 is postponed to Section 5 since it needs some preparation. It turns out that the proof of the null-critical case (given in [51]) is the most subtle part of

the proof of Theorem 1.3. It relies on the large time behaviors of the parabolic capacity potential and of the heat content that are studied in Section 3, and on an extension of Varadhan's lemma that is proved in Section 4 (see Lemma 4.1). Section 6 is devoted to some applications to Theorem 1.3. In particular, Corollary 6.7 seems to be new.

The next two sections deal with ratio limits. In Section 7, we discuss the existence of the strong ratio limit

$$\lim_{t \rightarrow \infty} \frac{k_P^M(x, y, t)}{k_P^M(x_0, x_0, t)}$$

(Davies' conjecture), and in Section 8 we deal with the conjecture that

$$(1.10) \quad \lim_{t \rightarrow \infty} \frac{k_{P_+}^M(x, y, t)}{k_{P_0}^M(x, y, t)} = 0 \quad \forall x, y \in M,$$

where P_+ and P_0 are respectively subcritical and critical operators in M . Finally, in Section 9 we discuss the equivalence of heat kernels of two subcritical elliptic operators (with the same λ_0) that agree outside a compact set. Unlike the analogous question concerning the equivalence of Green functions, the problem concerning the equivalence of heat kernels is still "terra incognita". Solving this problem will in particular enable us to study the stability of the large time behavior of the heat kernel under perturbations.

The survey is an expanded version of a talk given by the author at the conference "Mathematical Physics, Spectral Theory and Stochastic Analysis" held in Goslar, Germany, September 11–16, 2011, in honor of Professor Michael Demuth. We note that most of the results in sections 3–6 originally appeared in [48, 51], those of Section 7 appeared in [52], while those in sections 8–9 originally appeared in [24].

2. PRELIMINARIES

In this section we recall basic definitions and facts concerning the theory of nonnegative solutions of second-order linear elliptic and parabolic operators (for more details and proofs, see for example [54]).

Let P be a linear, second-order, elliptic operator defined in a non-compact, connected, C^3 -smooth Riemannian manifold M of dimension d . Here P is an elliptic operator with real and Hölder continuous coefficients which in any coordinate system $(U; x_1, \dots, x_d)$ has the form

$$(2.1) \quad P(x, \partial_x) = - \sum_{i,j=1}^d a_{ij}(x) \partial_i \partial_j + \sum_{i=1}^d b_i(x) \partial_i + c(x),$$

where $\partial_i = \partial/\partial x_i$. We assume that for every $x \in M$ the real quadratic form

$$(2.2) \quad \sum_{i,j=1}^d a_{ij}(x) \xi_i \xi_j, \quad \xi = (\xi_1, \dots, \xi_d) \in \mathbb{R}^d$$

is symmetric and positive definite.

Throughout the paper we always assume that $\lambda_0(P, M) \geq 0$. In this case we simply say that the operator P is *nonnegative in M* , and write $P \geq 0$ in M . So, $P \geq 0$ in M if and only if the equation $Pu = 0$ admits a global positive solution in M . We note that in the symmetric case, by the Agmon-Allegretto-Piepenbrink theory [1], the nonnegativity of P in M is equivalent to the nonnegativity of the associated quadratic form on $C_0^\infty(M)$.

We consider the parabolic operator L

$$(2.3) \quad Lu = u_t + Pu \quad \text{on } M \times (0, \infty),$$

and the corresponding homogeneous equation $Lu = 0$ in $M \times (0, \infty)$.

Remark 2.1. We confine ourselves to classical solutions. However, since the results and the proofs throughout the paper rely on standard elliptic and parabolic regularity, and basic properties and results of potential theory, the results of the paper are also valid for weak solutions in the case where the elliptic operator P is in divergence form

$$(2.4) \quad Pu = - \sum_{i=1}^d \partial_i \left[\sum_{j=1}^d a_{ij}(x) \partial_j u + u \tilde{b}_i(x) \right] + \sum_{i=1}^d b_i(x) \partial_i u + c(x)u,$$

with coefficients which satisfy standard local regularity assumptions (as for example in Section 1.1 of [44]). The results are also valid in the framework of strong solutions, where the strictly elliptic operator P is of the form (2.1) and has locally bounded coefficients; the proofs differ only in minor details from the proofs given here.

We write $\Omega_1 \Subset \Omega_2$ if Ω_2 is open, $\overline{\Omega_1}$ is compact and $\overline{\Omega_1} \subset \Omega_2$. Let $f, g \in C(\Omega)$ be nonnegative functions; we use the notation $f \asymp g$ on Ω if there exists a positive constant C such that

$$C^{-1}g(x) \leq f(x) \leq Cg(x) \quad \text{for all } x \in \Omega.$$

Let $\{M_j\}_{j=1}^\infty$ be an *exhaustion* of M , i.e. a sequence of smooth, relatively compact domains in M such that $M_1 \neq \emptyset$, $M_j \Subset M_{j+1}$ and $\bigcup_{j=1}^\infty M_j = M$. For every $j \geq 1$, we denote $M_j^* = M \setminus \text{cl}(M_j)$. Let $M_\infty = M \cup \{\infty\}$ be the one-point compactification of M . By a neighborhood of infinity in M we mean a neighborhood of ∞ in M_∞ , that is, a set of the form $M \setminus K$, where K is compact in M .

Assume that $P \geq 0$ in M . For every $j \geq 1$, consider the Dirichlet heat kernel $k_P^{M_j}(x, y, t)$ of the parabolic operator $L = \partial_t + P$ in M_j . So, for every continuous function f with a compact support in M , the function

$$u(x, t) := \int_{M_j} k_P^{M_j}(x, y, t) f(y) \, dy$$

solves the initial-Dirichlet boundary value problems

$$(2.5) \quad \begin{aligned} Lu &= 0 & \text{in } M_j \times (0, \infty), \\ u &= 0 & \text{on } \partial M_j \times (0, \infty), \\ u &= f & \text{on } M_j \times \{0\}. \end{aligned}$$

By the generalized maximum principle, $\{k_P^{M_j}(x, y, t)\}_{j=1}^\infty$ is an increasing sequence which converges to $k_P^M(x, y, t)$, the *positive minimal heat kernel* of the parabolic operator L in M .

The main properties of the positive minimal heat kernel are summarized in the following lemma.

Lemma 2.2. *Assume that $P \geq 0$ in M . The heat kernel $k_P^M(x, y, t)$ satisfies the following properties.*

(1) *Positivity:* $k_P^M(x, y, t) \geq 0$ for all $t \geq 0$ and $x, y \in M$.

(2) *For any $\lambda \leq \lambda_0$ we have*

$$(2.6) \quad k_{P-\lambda}^M(x, y, t) = e^{\lambda t} k_P^M(x, y, t).$$

(3) *The heat kernel satisfies the semigroup identity*

$$(2.7) \quad \int_{\Omega} k_P^M(x, z, t) k_P^M(z, y, \tau) \, dz = k_P^M(x, y, t + \tau) \quad \forall t, \tau > 0, \text{ and } x, y \in M.$$

(4) *Monotonicity:* If $M_1 \subset M$, and $V_1 \geq V_2$, then

$$k_P^{M_1} \leq k_P^M, \quad \text{and} \quad k_{P+V_1}^M \leq k_{P+V_2}^M.$$

(5) *For any fixed $y \in M$ (respectively $x \in M$), $k_P^M(x, y, t)$ solves the equation $u_t + P(x, \partial_x)u = 0$ (respectively $u_t + P^*(y, \partial_y)u = 0$) in $M \times (0, \infty)$.*

(6) *Skew product operators:* Let $M = M_1 \times M_2$ be a product of two manifolds, and denote $x = (x_1, x_2) \in M = M_1 \times M_2$. Consider a skew product elliptic operator of the form $P := P_1 \otimes I_2 + I_1 \otimes P_2$, where P_i is a second-order elliptic operator on M_i satisfying the

assumptions of the present paper, and I_i is the identity map on M_i , $i = 1, 2$. Then

$$(2.8) \quad k_P^M(x, y, t) = k_{P_1}^{M_1}(x_1, y_1, t) k_{P_2}^{M_2}(x_2, y_2, t).$$

- (7) *Eigenfunction expansion:* Suppose that P is a symmetric elliptic operator with (up to the boundary) smooth coefficients which is defined on a smooth bounded domain M . Let $\{\varphi_n\}_{n=0}^\infty$ be the complete orthonormal sequence of the (Dirichlet) eigenfunctions of P with the corresponding nondecreasing sequence of eigenvalues $\{\lambda_n\}_{n=0}^\infty$. Then the heat kernel of P in M has the eigenfunction expansion

$$(2.9) \quad k_P^M(x, y, t) = \sum_{n=0}^{\infty} e^{-\lambda_n t} \phi_n(x) \phi_n(y).$$

- (8) Let $v \in \mathcal{C}_P(M)$ and $v^* \in \mathcal{C}_{P^*}(M)$. Then we have [48, 49]

$$(2.10) \quad \int_M k_P^M(x, y, t) v(y) dy \leq v(x), \quad \text{and} \quad \int_M k_P^M(x, y, t) v^*(x) dx \leq v^*(y).$$

Moreover, (by the maximum principle) either

$$\int_M k_P^M(x, y, t) v(y) dy < v(x) \quad \forall (x, t) \in M \times (0, \infty),$$

or

$$\int_M k_P^M(x, y, t) v(y) dy = v(x) \quad \forall (x, t) \in M \times (0, \infty),$$

and in the latter case v is called an invariant solution of the operator P on M (see for example [21, 26, 48, 54]).

Remark 2.3. 1. If there exists $v \in \mathcal{C}_P(M)$ such that v is not invariant, then the positive Cauchy problem

$$Lu = 0, \quad u \geq 0 \quad \text{on } M \times (0, \infty), \quad u(x, 0) = 0 \quad x \in M$$

does not admit a unique solution.

2. An invariant solution is sometimes called *complete*. If the constant function is an invariant solution with respect to the heat operator, then one says that the heat operator *conserves probability*, and the corresponding diffusion process is said to be *stochastically complete* [27].

Definition 2.4. Let w be a positive solution of the equation $Pu = 0$ in $M \setminus K$, where $K \Subset M$. We say that w is a positive solution of the equation $Pu = 0$ of *minimal growth in a neighborhood of infinity* in M if for any $v \in C(\text{cl}(M_j^*))$ which is a positive supersolution of the equation $Pu = 0$ in M_j^* for some $j \geq 1$ large enough, and satisfies $w \leq v$ on ∂M_j^* , we have $w \leq v$ in M_j^* [1, 48, 54].

In the subcritical (respectively, critical) case, the Green function $G_P^M(\cdot, y)$ (respectively, the ground state φ) is a positive solution of the equation $Pu = 0$ of minimal growth in a neighborhood of infinity in M . Recall that if $\lambda < \lambda_0$, then $P - \lambda$ is subcritical in M . In particular, if P is critical in M , then $\lambda_0 = 0$.

Next, we recall the parabolic Harnack inequality. We denote by $Q(x_0, t_0, R, \tau)$ the parabolic box

$$Q(x_0, t_0, R, \theta) := \{(x, t) \in M \times \mathbb{R} \mid \rho(x, x_0) < R, t \in (t_0, t_0 + \theta R^2)\},$$

where ρ is the given Riemannian metric on M . We have:

Lemma 2.5 (Harnack inequality). *Let u be a nonnegative solution of the equation $Lu = 0$ in $Q(x_0, t_0, R, \theta)$, and assume that $\theta > 1$ and $0 < R < R_0$. Then*

$$(2.11) \quad u(x_0, t_0 + R^2) \leq Cu(x, t_0 + \theta R^2)$$

for every $\rho(x, x_0) < R/2$, where $C = C(L, d, R_0, \theta)$. Moreover, C varies within bounded bounds for all $\theta > 1$ such that $0 < \varepsilon \leq (\theta - 1)^{-1} \leq M$.

Let $v \in \mathcal{C}_P(M)$ and $v^* \in \mathcal{C}_{P^*}(M)$. Then the parabolic Harnack inequality (2.11), and (2.10) imply the following important estimate

$$(2.12) \quad k_P^M(x, y, t) \leq c_1(y)v(x), \quad \text{and} \quad k_P^M(x, y, t) \leq c_2(x)v^*(y)$$

for all $x, y \in M$ and $t > 1$ (see [48, (3.29–30)]). Recall that in the critical case, by uniqueness, v and v^* are (up to a multiplicative constant) the ground states φ and φ^* of P and P^* respectively. Moreover, it is known that the ground state φ (respectively, φ^*) is a positive invariant solution of the operator P (respectively, P^*) in M . So,

$$(2.13) \quad \int_M k_P^M(x, y, t) \varphi(y) dy = \varphi(x), \quad \text{and} \quad \int_M k_P^M(x, y, t) \varphi^*(x) dx = \varphi^*(y).$$

We distinguish between two types of criticality.

Definition 2.6. A critical operator P is said to be *positive-critical* in M if $\varphi^* \varphi \in L^1(M)$, and *null-critical* in M if $\varphi^* \varphi \notin L^1(M)$.

Remark 2.7. Let $\mathbf{1}$ be the constant function on M , taking at any point $x \in M$ the value 1. Suppose that $P\mathbf{1} = 0$. Then P is sub-critical (respectively, positive-critical, null-critical) in M if and only if the corresponding diffusion process is transient (respectively, positive-recurrent, null-recurrent). For a thorough discussion of the probabilistic interpretation of criticality theory, see [54].

In fact, in the critical case it is natural to use the well known (Doob) h -transform with $h = \varphi$, where φ is the ground state of P . So,

$$P^\varphi u := \frac{1}{\varphi} P(\varphi u) \quad \text{and therefore} \quad k_{P^\varphi}^M(x, y, t) = \frac{1}{\varphi(x)} k_P^M(x, y, t) \varphi(y).$$

Clearly, P^φ is an elliptic operator which satisfies all our assumptions. Note that P^φ is null-critical (respectively, positive-critical) if and only if P is null-critical (respectively, positive-critical), and the ground states of P^φ and $(P^\varphi)^*$ are $\mathbf{1}$ and $\varphi^* \varphi$, respectively. Moreover,

$$\lim_{t \rightarrow \infty} k_{P^\varphi}^M(x, y, t) = 0 \quad \text{if and only if} \quad \lim_{t \rightarrow \infty} k_P^M(x, y, t) = 0.$$

Therefore, in the critical case, we may assume that

$$(A) \quad P\mathbf{1} = 0, \text{ and } P \text{ is a critical operator in } M.$$

It is well known that on a general noncompact manifold M , the solution of the Cauchy problem for the parabolic equation $Lu = 0$ is not uniquely determined (see for example [30] and the references therein). On the other hand, under Assumption (A), there is a unique *minimal* solution of the Cauchy problem and of certain initial-boundary value problems for *bounded* initial and boundary conditions. More precisely,

Definition 2.8. Assume that $P\mathbf{1} = 0$. Let f be a *bounded* continuous function on M . By the *minimal solution* u of the Cauchy problem

$$\begin{aligned} Lu &= 0 & \text{in } M \times (0, \infty), \\ u &= f & \text{on } M \times \{0\}, \end{aligned}$$

we mean the function

$$(2.14) \quad u(x, t) := \int_M k_P^M(x, y, t) f(y) \, dy.$$

Note that (2.12) implies that u in (2.14) is well defined.

Definition 2.9. Assume that $P\mathbf{1} = 0$. Let $B \Subset M_1$ be a smooth bounded domain such that $B^* := M \setminus \text{cl}(B)$ is connected. Assume that f is a bounded continuous function on B^* , and g is a bounded

continuous function on $\partial B \times (0, \infty)$. By the *minimal solution* u of the initial-boundary value problem

$$(2.15) \quad \begin{aligned} Lu &= 0 && \text{in } B^* \times (0, \infty), \\ u &= g && \text{on } \partial B \times (0, \infty), \\ u &= f && \text{on } B^* \times \{0\}, \end{aligned}$$

we mean the limit of the solutions u_j of the following initial-boundary value problems

$$\begin{aligned} Lu &= 0 && \text{in } (B^* \cap M_j) \times (0, \infty), \\ u &= g && \text{on } \partial B \times (0, \infty), \\ u &= 0 && \text{on } \partial M_j \times (0, \infty), \\ u &= f && \text{on } (B^* \cap M_j) \times \{0\}. \end{aligned}$$

Remark 2.10. It can be easily checked that the sequence $\{u_j\}$ is indeed a converging sequence which converges to a solution of the initial-boundary value problem (2.15).

Next, we recall some results concerning the theory of positive solutions of *elliptic* equations that we shall need in the sequel.

The first result is a *Liouville comparison theorem* in the symmetric case.

Theorem 2.11 ([53]). *Let P_0 and P_1 be two symmetric operators defined on M of the form*

$$(2.16) \quad P_j u = -m_j^{-1} \operatorname{div}(m_j A_j \nabla u) + V_j u \quad j = 0, 1.$$

Assume that the following assumptions hold true.

- (i) *The operator P_0 is critical in M . Denote by $\varphi \in \mathcal{C}_{P_0}(M)$ its ground state.*
- (ii) *$P_1 \geq 0$ in M , and there exists a real function $\psi \in H_{\text{loc}}^1(M)$ such that $\psi_+ \neq 0$, and $P_1 \psi \leq 0$ in M , where $u_+(x) := \max\{0, u(x)\}$.*
- (iii) *The following matrix inequality holds*

$$(2.17) \quad (\psi_+)^2(x) m_1(x) A_1(x) \leq C \varphi^2(x) m_0(x) A_0(x) \quad \text{for a.e. } x \in M,$$

where $C > 0$ is a positive constant.

Then the operator P_1 is critical in M , and ψ is its ground state. In particular, $\dim \mathcal{C}_{P_1}(M) = 1$ and $\lambda_0(P_1, M) = 0$.

In the sequel we shall also need to use results concerning small and semismall perturbations of a subcritical elliptic operator. These notions were introduced in [46] and [43] respectively, and are closely related to the stability of $\mathcal{C}_P(\Omega)$ under perturbation by a potential V .

Definition 2.12. Let P be a subcritical operator in M , and let V be a real valued potential defined on M .

(i) We say that V is a *small perturbation* of P in M if

$$(2.18) \quad \lim_{j \rightarrow \infty} \left\{ \sup_{x, y \in M_j^*} \int_{M_j^*} \frac{G_P^M(x, z) |V(z)| G_P^M(z, y)}{G_P^M(x, y)} dz \right\} = 0.$$

(ii) V is a *semismall perturbation* of P in M if for some $x_0 \in M$ we have

$$(2.19) \quad \lim_{j \rightarrow \infty} \left\{ \sup_{y \in M_j^*} \int_{M_j^*} \frac{G_P^M(x_0, z) |V(z)| G_P^M(z, y)}{G_P^M(x_0, y)} dz \right\} = 0.$$

Recall that small perturbations are semismall [43]. For semismall perturbations we have

Theorem 2.13 ([43, 46, 47]). *Let P be a subcritical operator in M . Assume that $V = V_+ - V_-$ is a semismall perturbation of P^* in M satisfying $V_- \neq 0$, where $V_{\pm}(x) = \max\{0, \pm V(x)\}$.*

Then there exists $\alpha_0 > 0$ such that $P_{\alpha} := P + \alpha V$ is subcritical in M for all $0 \leq \alpha < \alpha_0$ and critical for $\alpha = \alpha_0$.

Moreover, let φ be the ground state of $P + \alpha_0 V$ and let y_0 be a fixed reference point in M_1 . Then for any $0 \leq \alpha < \alpha_0$

$$\varphi \asymp G_{P_{\alpha}}^M(\cdot, y_0) \quad \text{in } M_1^*,$$

where the equivalence constant depends on α .

3. CAPACITORY POTENTIAL AND HEAT CONTENT

Our first result concerning the large time behavior of positive solutions is given by the following simple lemma that does not distinguish between null-critical and positive-critical operators.

Lemma 3.1. *Assume that $P\mathbf{1} = 0$ and that P is critical in M . Let $B := B(x_0, \delta) \subset\subset M$ be the ball of radius δ centered at x_0 , and suppose that $B^* = M \setminus \text{cl}(B)$ is connected. Let w be **the heat content of B^*** , i.e. the minimal nonnegative solution of the following initial-boundary value problem*

$$(3.1) \quad \begin{aligned} Lu &= 0 & \text{in } B^* \times (0, \infty), \\ u &= 0 & \text{on } \partial B \times (0, \infty), \\ u &= 1 & \text{on } B^* \times \{0\}. \end{aligned}$$

Then w is a decreasing function of t , and $\lim_{t \rightarrow \infty} w(x, t) = 0$ locally uniformly in B^ .*

Proof. Clearly,

$$(3.2) \quad w(x, t) = \int_{B^*} k_P^{B^*}(x, y, t) dy < \int_M k_P^M(x, y, t) dy = 1.$$

It follows that $0 < w < 1$ in $B^* \times (0, \infty)$. Let $\varepsilon > 0$. By the semigroup identity and (3.2),

$$(3.3) \quad \begin{aligned} w(x, t + \varepsilon) &= \int_{B^*} k_P^{B^*}(x, y, t + \varepsilon) dy = \\ &= \int_{B^*} \left(\int_{B^*} k_P^{B^*}(x, z, t) k_P^{B^*}(z, y, \varepsilon) dz \right) dy = \end{aligned}$$

$$(3.4) \quad \begin{aligned} \int_{B^*} k_P^{B^*}(x, z, t) \left(\int_{B^*} k_P^{B^*}(z, y, \varepsilon) dy \right) dz &< \\ \int_{B^*} k_P^{B^*}(x, z, t) dz &= w(x, t). \end{aligned}$$

Hence, w is a decreasing function of t , and therefore, $\lim_{t \rightarrow \infty} w(x, t)$ exists. We denote $v(x) := \lim_{t \rightarrow \infty} w(x, t)$. Note that the above argument shows that even in the subcritical case w is a decreasing function of t .

For $\tau > 0$, consider the function $\nu(x, t; \tau) := w(x, t + \tau)$, where $t > -\tau$. Then $\nu(x, t; \tau)$ is a nonnegative solution of the parabolic equation $Lu = 0$ in $B^* \times (-\tau, \infty)$ which satisfies $u = 0$ on $\partial B \times (-\tau, \infty)$. By a standard parabolic argument as $\tau \rightarrow \infty$, any converging subsequence of this set of solutions converges locally uniformly to a solution of the parabolic equation $Lu = 0$ in $B^* \times \mathbb{R}$ which satisfies $u = 0$ on $\partial B \times \mathbb{R}$. Since $\lim_{\tau \rightarrow \infty} \nu(x, t; \tau) = \lim_{\tau \rightarrow \infty} w(x, \tau) = v(x)$, the limit does not depend on t , and v is a solution of the elliptic equation $Pu = 0$ in B^* , and satisfies $v = 0$ on ∂B . Furthermore, $0 \leq v < \mathbf{1}$.

Therefore, $\mathbf{1} - v$ is a positive solution of the equation $Pu = 0$ in B^* which satisfies $u = 1$ on ∂B . On the other hand, it follows from the criticality assumption that $\mathbf{1}$ is the minimal positive solution of the equation $Pu = 0$ in B^* which satisfies $u = 1$ on ∂B . Thus, $\mathbf{1} \leq \mathbf{1} - v$, and therefore, $v = 0$. \square

Definition 3.2. Let $B := B(x_0, \delta) \subset\subset M$. Suppose that $B^* = M \setminus \text{cl}(B)$ is connected. The nonnegative (minimal) solution

$$v(x, t) = \mathbf{1} - \int_{B^*} k_P^{B^*}(x, y, t) dy$$

is called the *parabolic capacity potential* of B^* . Note that v is indeed the minimal nonnegative solution of the initial-boundary value problem

$$(3.5) \quad \begin{aligned} Lu &= 0 && \text{in } B^* \times (0, \infty), \\ u &= 1 && \text{on } \partial B \times (0, \infty), \\ u &= 0 && \text{on } B^* \times \{0\}. \end{aligned}$$

Corollary 3.3. *Under the assumptions of Lemma 3.1, the parabolic capacity potential v of B^* is an increasing function of t , and we have $\lim_{t \rightarrow \infty} v(x, t) = 1$ locally uniformly in B^* .*

Proof. Using an exhaustion argument it is easily verified that

$$(3.6) \quad v(x, t) = \mathbf{1} - \int_{B^*} k_P^{B^*}(x, y, t) dy = \mathbf{1} - w(x, t),$$

where w is the heat content of B^* . Therefore, the corollary follows directly from Lemma 3.1. \square

4. VARADHAN'S LEMMA

Varadhan's celebrated lemma (see, [62, Lemma 9, p. 259] or [54, pp. 192–193]) deals with the large time behavior of minimal solutions of the Cauchy problem with *bounded* initial data (assuming that $P\mathbf{1} = 0$). It turns out that the limit as $t \rightarrow \infty$ of such solutions might not exist, but, under further conditions, the spacial oscillation of the solution tends to zero as $t \rightarrow \infty$. In the present section we slightly extend Varadhan's lemma (Lemma 4.1). This extended version of the lemma is crucially used in the proof of the null-critical case in Theorem 1.3.

Varadhan proved his lemma for *positive-critical* operators on \mathbb{R}^d using a purely probabilistic approach (ours is purely analytic). Our key observation is that the assertion of Varadhan's lemma is valid in our general setting under the weaker assumption that the skew product operator $\bar{P} := P \otimes I + I \otimes P$ is critical in $\bar{M} := M \times M$, where I is the identity operator on M . Note that if \bar{P} is critical in \bar{M} , then P is critical in M . On the other hand, if P is positive-critical in M , then \bar{P} is positive-critical in \bar{M} . Moreover, if \bar{P} is subcritical in \bar{M} , then by part (i) of Theorem 1.3, the heat kernel of \bar{P} on \bar{M} tends to zero as $t \rightarrow \infty$. Since the heat kernel of \bar{P} is equal to the product of the heat kernels of its factors (see Lemma 2.2), it follows that if \bar{P} is subcritical in \bar{M} , then $\lim_{t \rightarrow \infty} k_P^M(x, y, t) = 0$.

Consider the Riemannian product manifold $\bar{M} = M \times M$. A point in \bar{M} is denoted by $\bar{x} = (x_1, x_2)$. By P_{x_i} , $i = 1, 2$, we denote the operator P in the variable x_i . So, $\bar{P} = P_{x_1} + P_{x_2}$ is in fact the above skew

product operator defined on \bar{M} . We denote by \bar{L} the corresponding parabolic operator.

Lemma 4.1 (Varadhan's lemma [51]). *Assume that $P\mathbf{1} = 0$. Suppose further that \bar{P} is critical on \bar{M} . Let f be a continuous bounded function on M , and let*

$$u(x, t) = \int_M k_P^M(x, y, t) f(y) \, dy$$

be the minimal solution of the Cauchy problem with initial data f on M . Fix $K \subset\subset M$. Then

$$\lim_{t \rightarrow \infty} \sup_{x_1, x_2 \in K} |u(x_1, t) - u(x_2, t)| = 0.$$

Proof. Denote by $\bar{u}(\bar{x}, t) := u(x_1, t) - u(x_2, t)$. Recall that the heat kernel $\bar{k}(\bar{x}, \bar{y}, t)$ of the operator \bar{L} on \bar{M} satisfies

$$(4.1) \quad \bar{k}_{\bar{P}}^{\bar{M}}(\bar{x}, \bar{y}, t) = k_P^M(x_1, y_1, t) k_P^M(x_2, y_2, t).$$

By (2.13) and (4.1), we have

$$\begin{aligned} \bar{u}(\bar{x}, t) &= u(x_1, t) - u(x_2, t) = \\ &= \int_M k_P^M(x_1, y_1, t) f(y_1) \, dy_1 - \int_M k_P^M(x_2, y_2, t) f(y_2) \, dy_2 = \\ &= \int_M \int_M k_P^M(x_1, y_1, t) k_P^M(x_2, y_2, t) (f(y_1) - f(y_2)) \, dy_1 \, dy_2 = \\ &= \int_{\bar{M}} \bar{k}_{\bar{P}}^{\bar{M}}(\bar{x}, \bar{y}, t) (f(y_1) - f(y_2)) \, d\bar{y}. \end{aligned}$$

Hence, \bar{u} is the minimal solution of the Cauchy problem for the equation $\bar{L}\bar{u} = 0$ on $\bar{M} \times (0, \infty)$ with the bounded initial data $\bar{f}(\bar{x}) := f(x_1) - f(x_2)$, where $\bar{x} \in \bar{M}$.

Fix a compact set $K \subset\subset M$ and $x_0 \in M \setminus K$, and let $\varepsilon > 0$. Let $B := B((x_0, x_0), \delta) \subset\subset \bar{M} \setminus \bar{K}$, where $\bar{K} = K \times K$, and δ will be determined below. We may assume that $B^* = \bar{M} \setminus \text{cl}(B)$ is connected. Then \bar{u} is a minimal solution of the following initial-boundary value problem

$$(4.2) \quad \begin{aligned} \bar{L}\bar{u} &= 0 && \text{in } B^* \times (0, \infty), \\ \bar{u}(\bar{x}, t) &= u(x_1, t) - u(x_2, t) && \text{on } \partial B \times (0, \infty), \\ \bar{u}(\bar{x}, 0) &= f(x_1) - f(x_2) && \text{on } B^* \times \{0\}. \end{aligned}$$

We need to prove that $\lim_{t \rightarrow \infty} \bar{u}(\bar{x}, t) = 0$.

By the superposition principle (which obviously holds for minimal solutions), we have

$$\bar{u}(\bar{x}, t) = u_1(\bar{x}, t) + u_2(\bar{x}, t) \quad \text{on } B^* \times [1, \infty),$$

where u_1 solves the initial-boundary value problem

$$(4.3) \quad \begin{aligned} \bar{L}u_1 &= 0 && \text{in } B^* \times (1, \infty), \\ u_1(\bar{x}, t) &= u(x_1, t) - u(x_2, t) && \text{on } \partial B \times (1, \infty), \\ u_1(\bar{x}, 0) &= 0 && \text{on } B^* \times \{1\}, \end{aligned}$$

and u_2 solves the initial-boundary value problem

$$(4.4) \quad \begin{aligned} \bar{L}u_2 &= 0 && \text{in } B^* \times (1, \infty), \\ u_2(\bar{x}, t) &= 0 && \text{on } \partial B \times (1, \infty), \\ u_2(\bar{x}, 0) &= u(x_1, 1) - u(x_2, 1) && \text{on } B^* \times \{1\}. \end{aligned}$$

Clearly, $|\bar{u}(\bar{x}, t)| \leq 2\|f\|_\infty$ on $\bar{M} \times (0, \infty)$. Note that if $\bar{x} = (x_1, x_2) \in \partial B$, then on M , $\text{dist}_M(x_1, x_2) \leq 2\delta$. Using Schauder's parabolic interior estimates on M , it follows that if δ is small enough, then

$$|\bar{u}(\bar{x}, t)| = |u(x_1, t) - u(x_2, t)| < \varepsilon \quad \text{on } \partial B \times (1, \infty).$$

By comparison of u_1 with the parabolic capacity potential of B^* , we obtain that

$$(4.5) \quad |u_1(\bar{x}, t)| \leq \varepsilon \left(1 - \int_{B^*} \bar{k}_P^{B^*}(\bar{x}, \bar{y}, t-1) d\bar{y} \right) < \varepsilon \quad \text{in } B^* \times (1, \infty).$$

On the other hand,

$$(4.6) \quad |u_2(\bar{x}, t)| \leq 2\|f\|_\infty \int_{B^*} \bar{k}_P^{B^*}(\bar{x}, \bar{y}, t-1) d\bar{y} \quad \text{in } B^* \times (1, \infty).$$

It follows from (4.6) and Lemma 3.1 that there exists $T > 0$ such that

$$(4.7) \quad |u_2(\bar{x}, t)| \leq \varepsilon \quad \text{for all } \bar{x} \in \bar{K} \text{ and } t > T.$$

Combining (4.5) and (4.7), we obtain that $|u(x_1, t) - u(x_2, t)| \leq 2\varepsilon$ for all $x_1, x_2 \in K$ and $t > T$. Since ε is arbitrary, the lemma is proved. \square

5. EXISTENCE OF $\lim_{t \rightarrow \infty} e^{\lambda_0 t} k_P^M(x, y, t)$

The present section is devoted to the proof of Theorem 1.3 which claims that $\lim_{t \rightarrow \infty} e^{\lambda_0 t} k_P^M(x, y, t)$ always exists. Without loss of generality, we may assume that $\lambda_0 = 0$. For convenience, we denote by k the heat kernel k_P^M of the operator P in M .

Proof of Theorem 1.3. (i) *The subcritical case:* Suppose that P is subcritical in Ω . It means that for any $x, y \in M$ we have

$$(5.1) \quad \int_1^\infty k(x, y, t) dt < \infty.$$

Let $x, y \in \Omega$ be fixed, and suppose that $k(x, y, t)$ does not converge to zero as $t \rightarrow \infty$. Then there exist an increasing sequence $t_j \rightarrow \infty$,

$t_{j+1} - t_j > 1$, and $\varepsilon > 0$, such that $k(x, y, t_j) > \varepsilon$ for all $j > 1$. Using the parabolic Harnack inequality (2.11), we deduce that there exists $C > 0$ (C may depend on x, y, Ω , and P) such that

$$(5.2) \quad \int_{t_j}^{t_{j+1}} k(x, y, t) dt > C\varepsilon \quad \forall j > 2.$$

But this contradicts (5.1).

(ii) *The positive-critical case:* Let P be a critical operator in M and let $\{t_j\}_{j=1}^\infty \subset \mathbb{R}$ be a sequence such that $t_j \rightarrow \infty$, and define:

$$u_j(x, y, t) = k(x, y, t + t_j).$$

Fix $x_0, y_0 \in M$. By (2.12) we have

$$(5.3) \quad u_j(x, y_0, t) \leq c_1(y_0)\varphi(x), \quad \left(u_j(x_0, y, t) \leq c_2(x_0)\varphi^*(y), \right)$$

for all $x \in M$ ($y \in M$) and $t \in \mathbb{R}$ and $j > J(t)$. Using the parabolic Harnack inequality and a standard Parabolic regularity argument we may subtract a subsequence of $\{u_j\}$ (which we rename by $\{u_j\}$) that converges locally uniformly to a nonnegative solution $u(x, y, t)$ of the parabolic equations

$$Lu = \partial_t u + P(x, \partial_x)u = 0, \quad Lu = \partial_t u + P(y, \partial_y)u = 0 \quad \text{in } M \times \mathbb{R}.$$

Moreover, u satisfies the estimates

$$(5.4) \quad u(x, y_0, t) \leq c_1(y_0)\varphi(x) \quad \forall (x, t) \in M \times \mathbb{R},$$

$$(5.5) \quad u(x_0, y, t) \leq c_2(x_0)\varphi^*(y) \quad \forall (y, t) \in M \times \mathbb{R}.$$

Using the semigroup property we have for all $t, \tau > 0$

$$(5.6) \quad \int_M k(x, z, \tau)k(z, y, t + t_j) dz = \int_M k(x, z, t + t_j)k(z, y, \tau) dz$$

$$(5.7) \quad = k(x, y, \tau + t + t_j)$$

for all $j \geq 1$. On the other hand, by (2.13), the ground state φ is a positive invariant solution. Consequently, for any $x \in M$ and $\tau > 0$, $k(x, \cdot, \tau)\varphi \in L^1(M)$. Similarly, for all $\tau > 0$ and $y \in M$, $\varphi^*k(\cdot, y, \tau) \in L^1(M)$. Hence, by estimates (5.3) and the Lebesgue dominated convergence theorem, we obtain

$$(5.8) \quad \int_M k(x, z, \tau)u(z, y, t) dz = \int_M u(x, z, t)k(z, y, \tau) dz = u(x, y, \tau + t),$$

where $u(x, y, t) = \lim_{j \rightarrow \infty} k(x, y, t + t_j)$. In particular, for $\tau = t_j$ and $t = -t_j$ we have

$$(5.9) \quad \int_M k(x, z, t_j) u(z, y, -t_j) dz = \int_M u(x, z, -t_j) k(z, y, t_j) dz = u(x, y, 0)$$

Invoking again estimates (2.12), and (5.4)–(5.5), we see that the integrands in (5.9) are bounded by $C(x, y)\varphi(z)\varphi^*(z)$. Recall our assumption that $\varphi(x)\varphi^*(x) \in L^1(M)$, hence, the Lebesgue dominated convergence theorem implies

$$(5.10) \quad \int_M u(x, z, 0) \hat{u}(z, y, 0) dz = \int_M \hat{u}(x, z, 0) u(z, y, 0) dz = u(x, y, 0),$$

where

$$(5.11) \quad \hat{u}(x, y, t) := \lim_{j \rightarrow \infty} u(x, y, t - t_j),$$

and again we may assume that the limit in (5.11) exists.

On the other hand, by the invariance property (2.13), and estimates (2.12), we get

$$(5.12) \quad \int_M u(x, z, \tau) \varphi(z) dz = \varphi(x), \quad \int_M \varphi^*(z) u(z, y, t) dz = \varphi^*(y).$$

In particular $u(x, y, t) > 0$. It follows from the Harnack inequality and (5.10) that $\hat{u}(x, y, t)$ is also positive and we have

$$(5.13) \quad \int_M \int_M \hat{u}(x, z, 0) u(z, y, 0) dz \varphi(y) dy = \int_M u(x, y, 0) \varphi(y) dy.$$

Consequently, (5.12) and (5.13) imply that

$$(5.14) \quad \int_M \hat{u}(x, z, 0) \varphi(z) dz = \varphi(x).$$

Define integral operators

$$(5.15) \quad Uf(x) := \int_M \hat{u}(x, z, 0) f(z) dz, \quad U^*f(y) := \int_M \hat{u}(z, y, 0) f(z) dz.$$

By (5.10) and (5.14) we see that $\varphi(x)$ and $u(x, y, 0)$ (as a function of x) are positive eigenfunctions of the operator U with an eigenvalue 1, and for every $x \in M$, $u(x, y, 0)$ is a positive eigenfunction of U^* with an eigenvalue 1. Moreover, for every $x \in M$, $u(x, \cdot, 0)\varphi \in L^1(M)$ and for every $x, z \in M$, $u(x, \cdot, 0)u(\cdot, z, 0) \in L^1(M)$. Consequently, it follows

[48, Lemma 3.4] that 1 is a simple eigenvalue of the integral operator U . Hence,

$$(5.16) \quad u(x, y, 0) = \beta(y)\varphi(x).$$

By (5.12) and (5.16) we have

$$(5.17) \quad \int_M \varphi^*(z)\beta(y)\varphi(z) \, dz = \int_M \varphi^*(z)u(z, y, 0) \, dz = \varphi^*(y).$$

Therefore, we obtain from (5.16) and (5.17) that $\beta(y) = \frac{\varphi^*(y)}{\int_M \varphi(z)\varphi^*(z) \, dz}$.

Thus,

$$(5.18) \quad \lim_{j \rightarrow \infty} k(x, y, t_j) = u(x, y, 0) = \frac{\varphi(x)\varphi^*(y)}{\int_M \varphi(z)\varphi^*(z) \, dz},$$

and the positive-critical case is proved since the limit in (5.18) is independent of the sequence $\{t_j\}$.

(iii) *The null-critical case:* Without loss of generality, we may assume that $P\mathbf{1} = 0$, where P is a null-critical operator in M . We need to prove that

$$\lim_{t \rightarrow \infty} k_P^M(x, y, t) = 0.$$

As in Lemma 4.1 (Varadhan's lemma), consider the Riemannian product manifold $\bar{M} := M \times M$, and let $\bar{P} = P_{x_1} + P_{x_2}$ be the corresponding skew product operator which is defined on \bar{M} .

If \bar{P} is subcritical on \bar{M} , then by part (i), $\lim_{t \rightarrow \infty} \bar{k}_{\bar{P}}^{\bar{M}}(\bar{x}, \bar{y}, t) = 0$, and since

$$\bar{k}_{\bar{P}}^{\bar{M}}(\bar{x}, \bar{y}, t) = k_P^M(x_1, y_1, t)k_P^M(x_2, y_2, t),$$

it follows that $\lim_{t \rightarrow \infty} k_P^M(x, y, t) = 0$.

Therefore, there remains to prove the theorem for the case where \bar{P} is critical in \bar{M} . Fix a nonnegative, bounded, continuous function $f \neq 0$ such that $\varphi^*f \in L^1(M)$, and consider the solution

$$v(x, t) := \int_M k_P^M(x, y, t)f(y) \, dy.$$

Let $t_n \rightarrow \infty$, and consider the sequence $\{v_n(x, t) := v(x, t + t_n)\}$. As in part (ii), up to a subsequence, $\{v_n\}$ converges to a nonnegative solution $u \in \mathcal{H}_P(M \times \mathbb{R})$.

Invoking Lemma 4.1 (Varadhan's lemma), we see that $u(x, t) = \alpha(t)$. Since u solves the parabolic equation $Lu = 0$, it follows that $\alpha(t)$ is a nonnegative constant α .

We claim that $\alpha = 0$. Suppose to the contrary that $\alpha > 0$. The assumption that $\varphi^* f \in L^1(M)$ and (2.13) imply that for any $t > 0$

$$\begin{aligned} \int_M \varphi^*(y) v(y, t) dy &= \int_M \varphi^*(y) \left(\int_M k_P^M(y, z, t) f(z) dz \right) dy = \\ (5.19) \quad &\left(\int_M \varphi^*(y) k_P^M(y, z, t) dy \right) f(z) dz = \int_M \varphi^*(z) f(z) dz < \infty. \end{aligned}$$

On the other hand, by the null-criticality, Fatou's lemma, and (5.19) we have

$$\begin{aligned} \infty &= \int_M \varphi^*(z) \alpha dz = \int_M \varphi^*(z) \lim_{n \rightarrow \infty} v(z, t_n) dz \leq \\ \liminf_{n \rightarrow \infty} \int_M \varphi^*(z) v(z, t_n) dz &= \int_M \varphi^*(z) f(z) dz < \infty. \end{aligned}$$

Hence $\alpha = 0$, and therefore

$$(5.20) \quad \lim_{t \rightarrow \infty} \int_M k_P^M(x, y, t) f(y) dy = \lim_{t \rightarrow \infty} v(x, t) = 0.$$

Now fix $y \in M$ and let $f := k_P^M(\cdot, y, 1)$. Consider the minimal solution of the Cauchy problem with initial data f . So, by the semigroup property we have

$$\begin{aligned} u(x, t) &:= \int_M k_P^M(x, z, t) f(z) dz = \\ &\int_M k_P^M(x, z, t) k_P^M(z, y, 1) dz = k_P^M(x, y, t + 1). \end{aligned}$$

In view of (2.12) (with $v = \mathbf{1}$) and (2.13), the function f is bounded and satisfies $f\varphi^* \in L^1(M)$. Therefore, by (5.20) $\lim_{t \rightarrow \infty} u(x, t) = 0$. Thus,

$$\lim_{t \rightarrow \infty} k_P^M(x, y, t) = \lim_{t \rightarrow \infty} k_P^M(x, y, t + 1) = \lim_{t \rightarrow \infty} u(x, t) = 0.$$

The last statement of the theorem concerning the behavior of the Green function $G_{P-\lambda}^M(x, y)$ as $\lambda \rightarrow \lambda_0$ follows from the first part of the theorem using a classical Abelian theorem [58, Theorem 10.2]. \square

6. APPLICATIONS OF THEOREM 1.3

We discuss in this section some applications of Theorem 1.3 and comment on related results. First we prove (1.5), which characterizes λ_0 in terms of the large time behavior of $\log k_P^M(x, y, t)$.

Corollary 6.1 ([24]). *The heat kernel satisfies*

$$(6.1) \quad \lim_{t \rightarrow \infty} \frac{\log k_P^M(x, y, t)}{t} = -\lambda_0(P, M) \quad x, y \in M.$$

Proof. The needed upper bound for the validity of (6.1) follows directly from Theorem 1.3 and (1.4).

The lower bound is obtained by a standard exhaustion argument and Theorem 1.3 (cf. the proof of [27, Theorem 10.24]). Indeed, let $\{M_j\}_{j=1}^\infty$ be an *exhaustion* of M . Recall that since M_j is a smooth bounded domain, the operator $P - \lambda_0(P, M_j)$ is positive-critical in M_j . Therefore, Theorem 1.3, and the monotonicity of heat kernels with respect to domains (see part 4 of Lemma 2.2) imply that

$$\liminf_{t \rightarrow \infty} \frac{\log k_P^M(x, y, t)}{t} \geq \lim_{t \rightarrow \infty} \frac{\log k_P^{M_j}(x, y, t)}{t} = -\lambda_0(P, M_j).$$

Since $\lim_{j \rightarrow \infty} \lambda_0(P, M_j) = \lambda_0(P, M)$, we obtain the needed lower bound. \square

We now use Theorem 1.3 to strengthen Lemma 4.1. More precisely, in the following result we obtain the large time behavior of solutions of the Cauchy problem with initial conditions which satisfy a certain (and in some sense optimal) integrability condition.

Corollary 6.2 ([51]). *Let P be an elliptic operator of the form (2.1) such that $\lambda_0 \geq 0$. Let f be a continuous function on M such that $v^*f \in L^1(M)$ for some $v^* \in \mathcal{C}_{P^*}(M)$. Let*

$$u(x, t) = \int_M k_P^M(x, y, t) f(y) \, dy$$

be the minimal solution of the Cauchy problem with initial data f on M . Fix $K \subset\subset M$. Then

$$\lim_{t \rightarrow \infty} \sup_{x \in K} |u(x, t) - \mathcal{F}(x)| = 0,$$

where

$$\mathcal{F}(x) := \begin{cases} \frac{\int_M f(y) \varphi^*(y) \, dy}{\int_M \varphi(y) \varphi^*(y) \, dy} \varphi(x) & \text{if } P \text{ is positive-critical in } M, \\ 0 & \text{otherwise.} \end{cases}$$

Proof. Since $v^*f \in L^1(M)$, estimate (2.12), and the dominated convergence theorem imply that

$$\lim_{t \rightarrow \infty} u(x, t) = \lim_{t \rightarrow \infty} \int_M k_P^M(x, y, t) f(y) \, dy = \int_M \lim_{t \rightarrow \infty} [k_P^M(x, y, t) f(y)] \, dy,$$

and the claim of the corollary follows from Theorem 1.3. \square

Assume now that $P\mathbf{1} = 0$ and $\int_M k_P^M(\cdot, y, t) dy = \mathbf{1}$ (i.e. $\mathbf{1}$ is a positive invariant solution of the operator P in M). Corollary 6.2 implies that for any $j \geq 1$ and all $x \in M$ we have

$$\lim_{t \rightarrow \infty} \int_{M_j^*} k_P^M(x, y, t) dy = \begin{cases} \frac{\int_{M_j^*} \varphi^*(y) dy}{\int_M \varphi^*(y) dy} & \text{if } P \text{ is positive-critical in } M, \\ 1 & \text{otherwise.} \end{cases}$$

Suppose further that P is not positive-critical in M , and f is a bounded continuous function such that $\liminf_{x \rightarrow \infty} f(x) = \varepsilon > 0$. Then

$$(6.2) \quad \liminf_{t \rightarrow \infty} \int_M k_P^M(x, y, t) f(y) dy \geq \varepsilon.$$

Hence, if the integrability condition of Corollary 6.2 is not satisfied, then the large time behavior of the minimal solution of the Cauchy problem may be complicated. The following example of W. Kirsch and B. Simon [34] demonstrates this phenomenon.

Example 6.3. Consider the heat equation in \mathbb{R}^d . Let $R_j := e^{\varepsilon_j}$ and let

$$f(x) := 2 + (-1)^j \quad \text{if} \quad R_j < \sup_{1 \leq i \leq d} |x_i| < R_{j+1}, \quad j \geq 1.$$

Let u be the minimal solution of the Cauchy problem with initial data f . Then for $t \sim R_j R_{j+1}$ one has that $u(0, t) \sim 2 + (-1)^j$, and thus $u(0, t)$ does not have a limit. Note that by Lemma 4.1, for $d = 1$, $u(x, t)$ has exactly the same asymptotic behavior as $u(0, t)$ for all $x \in \mathbb{R}$.

In fact, it was proved in [56] that for the heat equation on \mathbb{R}^d , the following holds: for any bounded function f defined on \mathbb{R}^d , the limit

$$\lim_{t \rightarrow \infty} \int_{\mathbb{R}^d} k_{-\Delta}^{\mathbb{R}^d}(x, y, t) f(y) dy$$

exists, if and only if the limit

$$\lim_{R \rightarrow \infty} \frac{1}{|B(x, R)|} \int_{B(x, R)} f(y) dy$$

exists. Moreover, the values of the two limits are equal. For an extension of the above result see [31]. We note that such a theorem is false if the average value of f is taken on solid cubes.

The next three corollaries concern elliptic operators on manifolds.

Corollary 6.4 ([12]). *Let M be a noncompact complete Riemannian manifold, and denote by Δ the corresponding Laplace-Beltrami operator. Then*

$$(6.3) \quad \ell_0 := \lim_{t \rightarrow \infty} k_{-\Delta}^M(x, y, t) = 0$$

if and only if M has an infinite volume.

Proof. If M has a finite volume, then $-\Delta$ is positive-critical in M , and by Theorem 1.3, we have $\ell_0 = (\text{vol}(M))^{-1} > 0$.

Suppose now that M has an infinite volume. If $\lambda_0 > 0$, Corollary 6.1 implies that $\ell_0 = 0$. On the other hand, if $\lambda_0 = 0$, then since $u = \mathbf{1}$ is a positive harmonic function which is not in $L^2(M)$, the uniqueness of the ground state implies that $-\Delta$ is not positive-critical in M . Hence, by Theorem 1.3 we have $\ell_0 = 0$. \square

Before studying the next two results we recall three basic notions from the theory of manifolds with group actions.

Definition 6.5. Let G be a group, and suppose that G acts on M . For any real continuous function v and $g \in G$, denote by v^g the function defined by $v^g(x) := v(gx)$. Let $\mathbb{R}^* = \mathbb{R} \setminus \{0\}$ be the real multiplicative group.

1. A nonzero real continuous function f on M is called *G -multiplicative* if there exists a group homomorphism $\gamma : G \rightarrow \mathbb{R}^*$, such that

$$f(gx) = \gamma(g)f(x) \quad \forall g \in G, x \in M.$$

2. A G -group action on M is *compactly generating* if there exists $K \Subset M$ such that $GK = M$ (see [38]).

3. The operator P is said to be *G -equivariant* if

$$P[u^g] = (P[u])^g \quad \forall g \in G.$$

Corollary 6.6 (cf. [12]). *Suppose that a noncompact manifold M is a covering of a compact Riemannian manifold, and consider the Laplace-Beltrami operator Δ on M . Then*

$$(6.4) \quad \ell := \lim_{t \rightarrow \infty} e^{\lambda_0 t} k_{-\Delta}^M(x, y, t) = 0.$$

In particular, $-\Delta - \lambda_0$ is not positive-critical in M .

Proof. Suppose that $\ell > 0$. By Theorem 1.3, the operator $-\Delta - \lambda_0$ is positive-critical in M , and in particular, the ground state φ is in $L^2(M)$. The uniqueness of the ground state implies that φ is G -multiplicative, where G is the deck transformation. But this contradicts the assumption that $\varphi \in L^2$, since G is an infinite Group. \square

The following corollary deals with manifolds with group actions and *general* equivariant operators, without any assumption on the growth of the acting groups. It generalizes Corollary 6.6, and seems to be new.

Corollary 6.7. *Let M be a noncompact manifold, and let G be a group. Suppose that G acts on M , and that the G -group action on M is compactly generating. Assume that P is a G -equivariant second-order elliptic operator of the form (2.1) which is defined on M . Suppose further that any nonzero G -multiplicative $C^{2,\alpha}$ -function is not integrable on M . Then*

$$(6.5) \quad \ell := \lim_{t \rightarrow \infty} e^{\lambda_0 t} k_P^M(x, y, t) = 0.$$

Proof. Suppose that $\ell > 0$. By Theorem 1.3, the operator $P - \lambda_0$ is positive-critical in M . In particular, $\dim \mathcal{C}_{P-\lambda_0}(M) = \dim \mathcal{C}_{P^*-\lambda_0}(M) = 1$. By the uniqueness of the ground state, it follows that the corresponding ground states φ and φ^* are G -multiplicative. Therefore, their product $\varphi\varphi^*$ is also G -multiplicative, and we arrived at a contradiction since by our assumption a nonzero G -multiplicative $C^{2,\alpha}$ -function is not integrable on M . \square

Remark 6.8. 1. Corollary 6.7 clearly applies to the case where M is a regular covering of a compact Riemannian manifold, and G is the corresponding deck transformation. In fact, in this case, by [38, Theorem 5.17], the product of the ground states is a positive G -invariant function, and therefore it is not integrable on M .

2. Corollary 6.7 applies in particular to the case where P is defined on \mathbb{R}^d , and P is \mathbb{Z}^d -equivariant (that is, P has \mathbb{Z}^d -periodic coefficients). In this case it is known that $P - \lambda_0$ is (null)-critical if and only if $d = 1, 2$. For further related results, see [38].

Finally, let us introduce an important subclass of elliptic operators P such that $P - \lambda_0$ is positive-critical in M .

Definition 6.9. Assume that P is symmetric and $P - \lambda_0$ is positive-critical in M with a ground state φ satisfying the normalization condition $\|\varphi\|_{L^2(M)} = 1$. Let $T_P^M(t)$ be the corresponding (Dirichlet) semigroup on $L^2(M)$ generated by P . We say that $T_P^M(t)$ is *intrinsically ultracontractive* if for any $t > 0$ there exists a positive-constant C_t such that

$$0 \leq e^{\lambda_0 t} k_P^M(x, y, t) \leq c_t \varphi(x) \varphi(y) \quad \forall x, y \in M.$$

Example 6.10. If M is a smooth bounded domain and $P := -\Delta$, then the semigroup $T_P^M(t)$ is intrinsically ultracontractive on $L^2(M)$. Also, let $M := \mathbb{R}^d$ and $P := -\Delta + (1 + |x|^2)^{\alpha/2}$, where $\alpha \in \mathbb{R}$. Then

$T_P^M(t)$ is intrinsically ultracontractive on $L^2(M)$ if and only if $\alpha > 2$ (see also [20, 44] and references therein).

It turns out that in the intrinsically ultracontractive case the rate of the convergence (as $t \rightarrow \infty$) of the h -transformed heat kernel is uniformly exponentially fast.

Theorem 6.11 ([20, Theorem 4.2.5]). *Assume that P is symmetric and subcritical in M , and suppose that $T_P^M(t)$ is intrinsically ultracontractive on $L^2(M)$. Then there exists a complete orthonormal set $\{\varphi_j\}_{j=0}^\infty$ in $L^2(M)$ such that $\varphi_0 = \varphi$, and for any $j \geq 0$ the function φ_j is an eigenfunction of the Friedrichs extension of the operator P with eigenvalue λ_j , where $\{\lambda_j\}$ is nondecreasing. Moreover, the heat kernel has the eigenfunction expansion (2.9), and there exist positive-constants C and δ such that*

$$\left| e^{\lambda_0 t} \frac{k_P^M(x, y, t)}{\varphi_0(x)\varphi_0(y)} - 1 \right| \leq C e^{-\delta t} \quad \forall x, y \in M, t > 1.$$

Furthermore, for any $\varepsilon > 0$ the series

$$e^{\lambda_0 t} \frac{k_P^M(x, y, t)}{\varphi_0(x)\varphi_0(y)} = \sum_{n=0}^{\infty} e^{-\lambda_n t} \frac{\varphi_n(x)\varphi_n(y)}{\varphi_0(x)\varphi_0(y)}$$

converges uniformly on $M \times M \times [\varepsilon, \infty)$.

Remark 6.12. Murata [45] proved that if $T_P^M(t)$ is intrinsically ultracontractive on $L^2(M)$, then $\mathbf{1}$ is a small perturbation of P in M . In particular, for any $n \geq 0$ the function φ_n/φ_0 is bounded, and has a continuous extension up to the Martin boundary of M (with respect to P) [50]. On the other hand, an example of Bañuelos and Davis in [7] gives us a finite area domain $M \subset \mathbb{R}^2$ such that $\mathbf{1}$ is a small perturbation of the Laplacian in M , but the corresponding semigroup is not intrinsically ultracontractive.

Remark 6.13. In the null-recurrent case, the heat kernel may decay very slowly as $t \rightarrow \infty$, and one can construct a complete Riemannian manifold M such that all its Riemannian products $M^j, j \geq 1$ are null-recurrent with respect to the Laplace-Beltrami operator on M^j (see [19]).

7. DAVIES' CONJECTURE CONCERNING STRONG RATIO LIMIT

Having proved in Section 5 that $\lim_{t \rightarrow \infty} e^{\lambda_0 t} k_P^M(x, y, t)$ always exists, we next ask how fast this limit is approached. It is natural to conjecture that the limit is approached equally fast for different points $x, y \in M$.

Note that in the context of Markov chains, such an *(individual) strong ratio limit property* is in general not true [13]. The following conjecture was raised by E. B. Davies [22] in the selfadjoint case.

Conjecture 7.1 (Davies' conjecture). *Let $Lu = u_t + P(x, \partial_x)u$ be a parabolic operator which is defined on a noncompact Riemannian manifold M , and assume that $\lambda_0(P, M) \geq 0$. Fix a reference point $x_0 \in M$. Then*

$$(7.1) \quad \lim_{t \rightarrow \infty} \frac{k_P^M(x, y, t)}{k_P^M(x_0, x_0, t)} = a(x, y)$$

exists and is positive for all $x, y \in M$.

The aim of the present section is to discuss Conjecture 7.1 and closely related problems, and to obtain some results under minimal assumptions. Since, the conjecture does not depend on the value of λ_0 , we assume throughout the present section that $\lambda_0 = 0$.

Conjecture 7.1 was recently disproved by G. Kozma [35] in the *discrete* setting:

Theorem 7.2 (Kozma [35]). *There exists a connected graph G with bounded weights and two vertices $x, y \in G$ such that*

$$(7.2) \quad \lim_{n \rightarrow \infty} \frac{k(x, x, n)}{k(y, y, n)}$$

does not exist, where k is the heat kernel of the lazy random walk such that the walker, at every step, chooses with probability $1/2$ to stay in place, and with probability $1/2$ to move to one of the neighbors (with probability proportional to the given weights).

In addition, Kozma indicates in [35] how the construction might be carried out in the category of Laplace-Beltrami operators on manifolds with bounded geometry. As noted in [35], the bounded weights property in the graph's setting is the analogue to the bounded geometry property in the manifold setting, a property that in fact was not assumed in Davies' conjecture. We note that in Kozma's example the ratio $\frac{k(x, x, n)}{k(y, y, n)}$ is bounded between two constants that are independent of t .

Nevertheless, as we show below, there are many situations where Conjecture 7.1 holds true.

Remark 7.3. 1. Theorem 1.3 implies that Conjecture 7.1 holds true in the positive-critical case, therefore, we assume throughout the present section that P is *not positive-critical*.

2. We also note that if P is *symmetric*, then the conjecture holds if

$$(7.3) \quad \dim \mathcal{C}_{P-\lambda_0}(M) = 1,$$

(i.e. in the “Liouvillian” case) see [4, Corollary 2.7]. In particular, it holds true for *critical selfadjoint* operators. It also holds for the Laplace-Beltrami operator on a complete Riemannian manifold of dimension d with nonnegative Ricci curvature [22].

3. Recently Agmon [2] obtained the exact asymptotics (in (x, y, t)) of the heat kernel for a \mathbb{Z}^d -periodic (non-selfadjoint) operator P on \mathbb{R}^d , and for an equivariant operator P defined on *abelian* cocompact covering manifold M . In particular, it follows from Agmon’s results that Conjecture 7.1 holds true in these cases. Note that in these cases (and even in the case of *nilpotent* cocompact covering) it is known [38] that

$$(7.4) \quad \dim \mathcal{C}_{P-\lambda_0}(M) = \dim \mathcal{C}_{P^*-\lambda_0}(M) = 1.$$

4. For other particular cases where the conjecture holds true see [4, 10, 14, 15, 22, 65].

Remark 7.4. It would be interesting to prove Conjecture 7.1 at least under the assumption

$$(7.5) \quad \dim \mathcal{C}_{P-\lambda_0}(M) = \dim \mathcal{C}_{P^*-\lambda_0}(M) = 1,$$

which holds true in the critical case and in many important subcritical (Liouvillian) cases. For a probabilistic interpretation of Conjecture 7.1, see [4].

Remark 7.5. Let $t_n \rightarrow \infty$. By a standard parabolic argument, we may extract a subsequence $\{t_{n_k}\}$ such that for every $x, y \in M$ and $s < 0$

$$(7.6) \quad a(x, y, s) := \lim_{k \rightarrow \infty} \frac{k_P^M(x, y, s + t_{n_k})}{k_P^M(x_0, y_0, t_{n_k})}$$

exists. Moreover, $a(\cdot, y, \cdot) \in \mathcal{H}_P(M \times \mathbb{R}_-)$, and $a(x, \cdot, \cdot) \in \mathcal{H}_{P^*}(M \times \mathbb{R}_-)$. Note that in the selfadjoint case, we can extract a subsequence $\{t_{n_k}\}$ such that the limit function a satisfies $a(\cdot, y, \cdot) \in \mathcal{H}_P(M \times \mathbb{R})$ [52].

Remark 7.6. Consider the complete (two-dimensional) Riemannian manifold M that is constructed in [49]. Then M does not admit non-constant positive harmonic functions, $\lambda_0(-\Delta, M) = 0$. Nevertheless, the heat operator does not admit any λ_0 -invariant positive solution. In particular, M is stochastically incomplete (this construction disproves

Stroock's conjecture concerning the existence of a λ_0 -invariant positive solution).

On the other hand, since M is Liouvillean, Remark 7.3 implies that Conjecture 7.1 holds true on M . Hence, the limit function

$$(7.7) \quad a(x, y) := \lim_{t \rightarrow \infty} \frac{k_{-\Delta}^M(x, y, t)}{k_{-\Delta}^M(x_0, x_0, t)}$$

(which equals to the constant function $\mathbf{1}$) is not a λ_0 -invariant positive solution. Compare this with [22, Theorem 25] and the discussion therein above Lemma 26. In particular, it follows that the function

$$\mu(x) := \sup \left\{ \sqrt{\frac{K_P^M(x, x, t)}{K_P^M(x_0, x_0, t)}} \mid 1 \leq t < \infty \right\}$$

is not slowly increasing in the sense of [22].

Suppose that P is symmetric (and $\lambda_o = 0$). Using (2.9) and a standard exhaustion argument, it follows [22] that for a fixed $x \in M$, the function $t \mapsto k_P^M(x, x, t)$ is a nonincreasing log-convex function, and therefore, a polarization argument implies that the following strong ratio property holds true.

$$(7.8) \quad \lim_{t \rightarrow \infty} \frac{k_P^M(x, y, t + s)}{k_P^M(x, y, t)} = 1 \quad \forall x, y \in M, s \in \mathbb{R}.$$

In the nonsymmetric case, Corollary 6.1 and the parabolic Harnack inequality imply:

Lemma 7.7 ([52]). *For every $x, y \in M$ and $s \in \mathbb{R}$, we have that*

$$(7.9) \quad \liminf_{t \rightarrow \infty} \frac{k_P^M(x, y, t + s)}{k_P^M(x, y, t)} \leq 1 \leq \limsup_{t \rightarrow \infty} \frac{k_P^M(x, y, t + s)}{k_P^M(x, y, t)}.$$

In particular, if $\lim_{t \rightarrow \infty} [k_P^M(x, y, t + s)/k_P^M(x, y, t)]$ exists, it equals to 1.

Remark 7.8. If there exist $x_0, y_0 \in M$ and $0 < s_0 < 1$ such that

$$(7.10) \quad M(x_0, y_0, s_0) := \limsup_{t \rightarrow \infty} \frac{k_P^M(x_0, y_0, t + s_0)}{k_P^M(x_0, y_0, t)} < \infty,$$

then by the parabolic Harnack inequality, for all $x, y, z, w \in K \subset\subset M$, $t > 1$, we have the following Harnack inequality of elliptic type:

$$(7.11) \quad k_P^M(z, w, t) \leq C_1 k_P^M(x_0, y_0, t + \frac{s_0}{2}) \leq C_2 k_P^M(x_0, y_0, t - \frac{s_0}{2}) \leq C_3 k_P^M(x, y, t).$$

Similarly, (7.10) implies that for all $x, y \in M$ and $r \in \mathbb{R}$:

$$0 < m(x, y, r) := \liminf_{t \rightarrow \infty} \frac{k_P^M(x, y, t + r)}{k_P^M(x_0, y_0, t)} \leq \limsup_{t \rightarrow \infty} \frac{k_P^M(x, y, t + r)}{k_P^M(x_0, y_0, t)} = M(x, y, r) < \infty.$$

Note that (7.10) is obviously satisfied in the symmetric case, and consequently (7.11) holds true [20, Theorem 10].

It turns out that the validity of the strong limit property (7.8) (in the nonsymmetric case) implies the validity of Davies' conjecture if in addition (7.5) is satisfied. We have:

Lemma 7.9 ([52]). *(a) The following assertions are equivalent:*

(i) For each $x, y \in M$ there exists a sequence $s_j \rightarrow 0$ of negative numbers such that for all $j \geq 1$, and $s = s_j$, we have

$$(7.12) \quad \lim_{t \rightarrow \infty} \frac{k_P^M(x, y, t + s)}{k_P^M(x, y, t)} = 1.$$

(ii) The ratio limit in (7.12) exists for any $x, y \in M$ and $s \in \mathbb{R}$.

(iii) Any limit function $u(x, y, s)$ of the quotients $\frac{k_P^M(x, y, t_n + s)}{k_P^M(x_0, y_0, t_n)}$ with $t_n \rightarrow \infty$ does not depend on s and has the form $u(x, y)$, where $u(\cdot, y) \in \mathcal{C}_P(M)$ for every $y \in M$ and $u(x, \cdot) \in \mathcal{C}_{P^}(M)$ for every $x \in M$.*

(b) If one assumes further that (7.5) is satisfied, then Conjecture 7.1 holds true.

Moreover, Conjecture 7.1 holds true if $M \subsetneq \mathbb{R}^d$ is a smooth unbounded domain, P and P^* are (up to the boundary) smooth operators, (7.12) holds true, and

$$(7.13) \quad \dim \mathcal{C}_P^0(M) = \dim \mathcal{C}_{P^*}^0(M) = 1,$$

where $\mathcal{C}_P^0(M)$ denotes the cone of all functions in $\mathcal{C}_P(M)$ that vanish on ∂M .

Proof. (a) By Lemma 7.7, if the limit in (7.12) exists, then it is 1.

(i) \Rightarrow (ii). Fix $x_0, y_0 \in M$, and take $s_0 < 0$ for which the limit (7.12) exists. It follows that any limit solution $u(x, y, s) \in \mathcal{H}_P(M \times \mathbb{R}_-)$ of a sequence $\left\{ \frac{k_P^M(x, y, t_n + s)}{k_P^M(x_0, y_0, t_n)} \right\}$ with $t_n \rightarrow \infty$ satisfies $u(x_0, y_0, s + s_0) = u(x_0, y_0, s)$ for all $s < 0$. So, $u(x_0, y_0, \cdot)$ is s_0 -periodic. It follows from our assumption and the continuity of u that $u(x_0, y_0, \cdot)$ is the constant

function. Since this holds for all $x, y \in M$ and u , it follows that (7.12) holds for any $x, y \in M$ and $s \in \mathbb{R}$.

(ii) \Rightarrow (iii). Fix $y \in M$. By Remark 7.5, any limit function u of the sequence $\left\{ \frac{k_P^M(x, y, t_n + s)}{k_P^M(x_0, x_0, t_n)} \right\}$ with $t_n \rightarrow \infty$ belongs to $\mathcal{H}_P(M \times \mathbb{R}_-)$. Since

$$(7.14) \quad \frac{k_P^M(x, y, t + s)}{k_P^M(x_0, x_0, t)} = \frac{k_P^M(x, y, t)}{k_P^M(x_0, x_0, t)} \frac{k_P^M(x, y, t + s)}{k_P^M(x, y, t)},$$

(7.12) implies that such a u does not depend on s . Therefore, $u = u(x, y)$, where $u(\cdot, y) \in \mathcal{C}_P(M)$ and $u(x, \cdot) \in \mathcal{C}_{P^*}(M)$.

(iii) \Rightarrow (i). Write

$$(7.15) \quad \frac{k_P^M(x, y, t + s)}{k_P^M(x, y, t)} = \frac{k_P^M(x, y, t + s)}{k_P^M(x_0, x_0, t)} \frac{k_P^M(x_0, x_0, t)}{k_P^M(x, y, t)}.$$

Let $t_n \rightarrow \infty$ be a sequence such that the sequence $\left\{ \frac{k_P^M(x, y, t_n + s)}{k_P^M(x_0, x_0, t_n)} \right\}$ converges to a solution in $\mathcal{H}_P(M \times \mathbb{R}_-)$. By our assumption, we have

$$\lim_{n \rightarrow \infty} \frac{k_P^M(x, y, t_n + s)}{k_P^M(x_0, x_0, t_n)} = \lim_{n \rightarrow \infty} \frac{k_P^M(x, y, t_n)}{k_P^M(x_0, x_0, t_n)} = u(x, y) > 0,$$

which together with (7.15) implies (7.12) for all $s \in \mathbb{R}$.

(b) The uniqueness and (iii) imply that $\frac{k_P^M(x, y, t + s)}{k_P^M(x_0, x_0, t)} \rightarrow \frac{u(x)u^*(y)}{u(x_0)u^*(x_0)}$, where $u \in \mathcal{C}_P(M)$ and $u^* \in \mathcal{C}_{P^*}(M)$, and Conjecture 7.1 holds. \square

Remark 7.10. Assume that one of the assumptions of part (a) of Lemma 7.9 is satisfied. Then by the lemma, any limit function of the sequence $\left\{ \frac{k_P^M(x, y, t_n + s)}{k_P^M(x_0, x_0, t_n)} \right\}$ is of the form $a(x, y)$, where for every $y \in M$, the function $a(\cdot, y) \in \mathcal{C}_P(M)$, and $a(x, \cdot) \in \mathcal{C}_{P^*}(M)$ for every $x \in M$. But in general, $a(x, y)$ does not need to be a *product* of solutions of the equations $Pu = 0$ and $P^*u = 0$, as is demonstrated in [15], in the hyperbolic space, and in [52, Example 4.2].

In the null-critical case we have:

Lemma 7.11 ([52]). *Suppose that P is null-critical, and for each $x, y \in M$ there exists a sequence $\{s_j\}$ of negative numbers such that $s_j \rightarrow 0$, and*

$$(7.16) \quad \liminf_{t \rightarrow \infty} \frac{k_P^M(x, y, t + s)}{k_P^M(x, y, t)} \geq 1$$

for $s = s_j$, $j = 1, 2, \dots$. Then Conjecture 7.1 holds true.

Proof. Let $u(x, y, s)$ be a limit function of a sequence $\left\{ \frac{k_P^{\mathcal{M}}(x, y, t_n + s)}{k_P^{\mathcal{M}}(x_0, x_0, t_n)} \right\}$ with $t_n \rightarrow \infty$ and $s < 0$. By our assumption, $u(x, y, s + s_j) \geq u(x, y, s)$, and therefore, $u_s(x, y, s) \leq 0$ for all $s < 0$. Thus, $u(\cdot, y, s)$ (respect., $u(x, \cdot, s)$) is a positive supersolution of the equation $Pu = 0$ (respect., $P^*u = 0$) in \mathcal{M} . Since P is critical, it follows that $u(\cdot, y, s) \in \mathcal{C}_P(\mathcal{M})$ (respect., $u(x, \cdot, s) \in \mathcal{C}_{P^*}(\mathcal{M})$), and hence $u_s(x, y, s) = 0$. By the uniqueness, u equals to $\frac{\varphi(x)\varphi^*(y)}{\varphi(x_0)\varphi^*(x_0)}$, and Conjecture 7.1 holds true. \square

The large time behavior of quotients of the heat kernel is obviously closely related to the parabolic Martin boundary (for the parabolic Martin boundary theory see [16, 23, 45]). The next result relates Conjecture 7.1 and the parabolic Martin compactification of $\mathcal{H}_P(M \times \mathbb{R}_-)$.

Theorem 7.12 ([52]). *Assume that (7.10) holds true for some $x_0, y_0 \in M$, and $s_0 > 0$. Then the following assertions are equivalent:*

(i) *Conjecture 7.1 holds true for a fixed $x_0 \in M$.*

(ii)

$$(7.17) \quad \lim_{t \rightarrow \infty} \frac{k_P^M(x, y, t)}{k_P^M(x_1, y_1, t)}$$

exists, and the limit is positive for every $x, y, x_1, y_1 \in M$.

(iii)

$$(7.18) \quad \lim_{t \rightarrow \infty} \frac{k_P^M(x, y, t)}{k_P^M(y, y, t)}, \quad \text{and} \quad \lim_{t \rightarrow \infty} \frac{k_P^M(x, y, t)}{k_P^M(x, x, t)}$$

exist, and these ratio limits are positive for every $x, y \in M$.

(iv) *For any $y \in M$ there is a unique nonzero parabolic Martin boundary point \bar{y} for the equation $Lu = 0$ in $M \times \mathbb{R}$ which corresponds to any sequence of the form $\{(y, -t_n)\}_{n=1}^\infty$ such that $t_n \rightarrow \infty$, and for any $x \in M$ there is a unique nonzero parabolic Martin boundary point \bar{x} for the equation $u_t + P^*u = 0$ in $M \times \mathbb{R}$ which corresponds to any sequence of the form $\{(x, -t_n)\}_{n=1}^\infty$ such that $t_n \rightarrow \infty$.*

Moreover, if Conjecture 7.1 holds true, then for any fixed $y \in M$ (respect., $x \in M$), the limit function $a(\cdot, y)$ (respect., $a(x, \cdot)$) is a positive solution of the equation $Pu = 0$ (respect., $P^*u = 0$). Furthermore, the Martin functions of part (iv) are time independent, and (7.12) holds for all $x, y \in M$ and $s \in \mathbb{R}$.

Proof. (i) \Rightarrow (ii) follows from the identity

$$\frac{k_P^M(x, y, t)}{k_P^M(x_1, y_1, t)} = \frac{k_P^M(x, y, t)}{k_P^M(x_0, x_0, t)} \cdot \left(\frac{k_P^M(x_1, y_1, t)}{k_P^M(x_0, x_0, t)} \right)^{-1}.$$

(ii) \Rightarrow (iii). Take $x_1 = y_1 = y$ and $x_1 = y_1 = x$, respectively.

(iii) \Rightarrow (iv). It is well known that the Martin compactification does not depend on the fixed reference point x_0 . So, fix $y \in M$ and take it also as a reference point. Let $\{-t_n\}$ be a sequence such that $t_n \rightarrow \infty$ and such that the Martin sequence $\left\{ \frac{k_P^M(x, y, t+t_n)}{k_P^M(y, y, t+t_n)} \right\}$ converges to a Martin function $K_P^M(x, \bar{y}, t)$. By our assumption, for any t we have

$$\lim_{n \rightarrow \infty} \frac{k_P^M(x, y, t+t_n)}{k_P^M(y, y, t+t_n)} = \lim_{\tau \rightarrow \infty} \frac{k_P^M(x, y, \tau)}{k_P^M(y, y, \tau)} = b(x) > 0,$$

where b does not depend on the sequence $\{-t_n\}$. On the other hand,

$$\lim_{n \rightarrow \infty} \frac{k_P^M(y, y, t+t_n)}{k_P^M(y, y, t_n)} = K_P^M(y, \bar{y}, t) = f(t).$$

Since

$$\frac{k_P^M(x, y, t+t_n)}{k_P^M(y, y, t_n)} = \frac{k_P^M(x, y, t+t_n)}{k_P^M(y, y, t+t_n)} \cdot \frac{k_P^M(y, y, t+t_n)}{k_P^M(y, y, t_n)},$$

we have

$$K_P^M(x, \bar{y}, t) = b(x)f(t).$$

By separation of variables, there exists a constant λ such that

$$Pb - \lambda b = 0 \quad \text{on } M, \quad f' + \lambda f = 0 \quad \text{on } \mathbb{R}, \quad f(0) = 1.$$

Since b does not depend on the sequence $\{-t_n\}$, it follows in particular, that λ does not depend on this sequence. Thus, $\lim_{\tau \rightarrow \infty} \frac{k_P^M(y, y, t+\tau)}{k_P^M(y, y, \tau)} = f(t) = e^{-\lambda t}$. Lemma 7.7 implies that $\lambda = 0$. It follows that b is a positive solution of the equation $Pu = 0$, and

$$(7.19) \quad K_P^M(x, \bar{y}, t) = \lim_{\tau \rightarrow -\infty} \frac{k_P^M(x, y, t-\tau)}{k_P^M(y, y, -\tau)} = b(x).$$

The dual assertion can be proved similarly.

(iv) \Rightarrow (i). Let $K_P^M(x, \bar{y}, t)$ be the Martin function given in (iv), and $s_0 > 0$ such that $K_P^M(x_0, \bar{y}, s_0/2) > 0$. Consequently, $K_P^M(x, \bar{y}, s) > 0$

for $s \geq s_0$. Using the substitution $\tau = s + s_0$ we obtain

$$(7.20) \quad \lim_{\tau \rightarrow \infty} \frac{k_P^M(x, y, \tau)}{k_P^M(x_0, x_0, \tau)} = \lim_{s \rightarrow \infty} \left\{ \frac{k_P^M(x, y, s + s_0)}{k_P^M(y, y, s)} \times \frac{k_P^M(y, y, s)}{k_P^M(x_0, y, s + 2s_0)} \frac{k_P^M(x_0, y, s + 2s_0)}{k_P^M(x_0, x_0, s + s_0)} \right\} = \frac{K_P^M(x, \bar{y}, s_0) K_{P^*}^M(\bar{x}_0, y, s_0)}{K_P^M(x_0, \bar{y}, 2s_0)}.$$

The last assertion of the theorem follows from (7.19) and Lemma 7.9. \square

Finally we mention a problem which was raised by Burdzy and Salisbury [9] for $P = -\Delta$ and $\mathcal{M} \subset \mathbb{R}^d$.

Question 7.13 ([52]). *Assume that $\lambda_0 = 0$. Determine which minimal functions (in the sense of Martin's boundary) in $\mathcal{C}_P(\mathcal{M})$ are minimal in $\mathcal{H}_P(\mathcal{M} \times \mathbb{R}_-)$. In particular, is it true that in the critical case, the ground state φ is minimal in $\mathcal{H}_P(\mathcal{M} \times \mathbb{R}_-)$?*

8. COMPARING DECAY OF CRITICAL AND SUBCRITICAL HEAT KERNELS

In this section we are concerned with the large time behavior of the heat kernel k_P^M with regards to the criticality versus subcriticality property of the operator P . Since for any fixed $x, y \in M$, $x \neq y$, we have that $k_P^M(x, y, \cdot) \in L^1(\mathbb{R}_+)$ if and only if P is subcritical, it is natural to conjecture that *under some assumptions* the heat kernel of a subcritical operator P_+ in M decays (in time) faster than the heat kernel of a critical operator P_0 in M . Hence, our aim is to discuss the following conjecture in *general* settings.

Conjecture 8.1 ([24]). *Let P_+ and P_0 be respectively subcritical and critical operators in M . Then*

$$(8.1) \quad \lim_{t \rightarrow \infty} \frac{k_{P_+}^M(x, y, t)}{k_{P_0}^M(x, y, t)} = 0$$

locally uniformly in $M \times M$.

Conjecture 8.1 was stimulated by the following conjecture of D. Krejčířík and E. Zuazua [36]:

Let P_+ and P_0 be, respectively, selfadjoint subcritical and critical operators defined on $L^2(M, dx)$. Then

$$(8.2) \quad \lim_{t \rightarrow \infty} \frac{\|e^{-P_+ t}\|_{L^2(M, W dx) \rightarrow L^2(M, dx)}}{\|e^{-P_0 t}\|_{L^2(M, W dx) \rightarrow L^2(M, dx)}} = 0$$

for some positive weight function W .

Remark 8.2. Theorem 1.3 implies that Conjecture 8.1 obviously holds true if P_0 is positive-critical. Moreover, by Corollary 6.1 the conjecture also holds true if $\lambda_0(P_+, M) > 0$. Therefore, throughout this section we assume that $\lambda_0(P_+, M) = 0$, and P_0 is null-critical in M .

Example 8.3. In [41] M. Murata obtained the exact asymptotic for the heat kernels of nonnegative Schrödinger operators with *short-range* (real) potentials defined on \mathbb{R}^d , $d \geq 1$. In particular, [41, theorems 4.2 and 4.4] imply that Conjecture 8.1 holds true for such operators.

The following theorem deals with the *symmetric* case.

Theorem 8.4 ([24]). *Let the subcritical operator P_+ and the critical operator P_0 be symmetric in M of the form*

$$(8.3) \quad Pu = -m^{-1} \operatorname{div}(mA \nabla u) + Vu.$$

Assume that A_+ and A_0 , the sections on M of $\operatorname{End}(TM)$, and the weights m_+ and m_0 , corresponding to P_+ and P_0 , respectively, satisfy the following matrix inequality

$$(8.4) \quad m_+(x)A_+(x) \leq Cm_0(x)A_0(x) \quad \text{for a.e. } x \in M,$$

where C is a positive constant. Assume further that for some fixed $y_1 \in M$ there exists a positive constant C satisfying the following condition: for each $x \in M$ there exists $T(x) > 0$ such that

$$(8.5) \quad k_{P_+}^M(x, y_1, t) \leq Ck_{P_0}^M(x, y_1, t) \quad \forall t > T(x).$$

Then

$$(8.6) \quad \lim_{t \rightarrow \infty} \frac{k_{P_+}^M(x, y, t)}{k_{P_0}^M(x, y, t)} = 0$$

locally uniformly in $M \times M$.

Proof. Recall that in light of Remark 8.2 we assume that $\lambda_0(P_+, M) = 0$. Suppose to the contrary that for some $x_0, y_0 \in M$ there exists a sequence $\{t_n\}$ such that $t_n \rightarrow \infty$, and

$$(8.7) \quad \lim_{n \rightarrow \infty} \frac{k_{P_+}^M(x_0, y_0, t_n)}{k_{P_0}^M(x_0, y_0, t_n)} = K > 0.$$

Consider the sequence of functions $\{u_n\}_{n=1}^\infty$ defined by

$$u_n(x, s) := \frac{k_{P_+}^M(x, y_0, t_n + s)}{k_{P_0}^M(x_0, y_0, t_n)} \quad x \in M, s \in \mathbb{R}.$$

We note that

$$u_n(x, s) = \frac{k_{P_+}^M(x, y_0, t_n + s)}{k_{P_+}^M(x_0, y_0, t_n)} \times \frac{k_{P_+}^M(x_0, y_0, t_n)}{k_{P_0}^M(x_0, y_0, t_n)}.$$

Therefore, by assumption (8.7) and Remark 7.5 it follows that we may subtract a subsequence which we rename by $\{u_n\}$ such that

$$\lim_{n \rightarrow \infty} u_n(x, s) = u_+(x, s),$$

where $u_+ \in \mathcal{H}_{P_+}(M \times \mathbb{R})$ and $u_+ \geq 0$.

On the other hand,

$$v_n(x) := \frac{k_{P_+}^M(x, y_0, t_n)}{k_{P_0}^M(x_0, y_0, t_n)} = u_n(x, s) \frac{k_{P_+}^M(x, y_0, t_n)}{k_{P_+}^M(x, y_0, t_n + s)}.$$

By our assumption, $\lambda_0(P_+, M) = 0$ and P_+ is symmetric, therefore (7.8) implies that

$$\lim_{n \rightarrow \infty} \frac{k_{P_+}^M(x, y_0, t_n)}{k_{P_+}^M(x, y_0, t_n + s)} = 1.$$

Therefore,

$$\lim_{n \rightarrow \infty} v_n(x) = \lim_{n \rightarrow \infty} u_n(x, s) = u_+(x, s),$$

and u_+ does not depend on s , and hence u_+ is a positive solution of the elliptic equation $P_+ u = 0$ in M and we have

$$(8.8) \quad \lim_{n \rightarrow \infty} \frac{k_{P_+}^M(x, y_0, t_n)}{k_{P_0}^M(x_0, y_0, t_n)} = u_+(x).$$

On the other hand, by Remark 7.3 we have

$$(8.9) \quad \lim_{n \rightarrow \infty} \frac{k_{P_0}^M(x, y_0, t_n)}{k_{P_0}^M(x_0, y_0, t_n)} = \frac{\varphi(x)}{\varphi(x_0)} =: u_0(x),$$

where φ is the ground state of P_0 .

Combining (8.8) and (8.9), we obtain

$$(8.10) \quad \lim_{n \rightarrow \infty} \frac{k_{P_+}^M(x, y_0, t_n)}{k_{P_0}^M(x, y_0, t_n)} = \lim_{n \rightarrow \infty} \left\{ \frac{\frac{k_{P_+}^M(x, y_0, t_n)}{k_{P_0}^M(x_0, y_0, t_n)}}{\frac{k_{P_0}^M(x, y_0, t_n)}{k_{P_0}^M(x_0, y_0, t_n)}} \right\} = \frac{u_+(x)}{u_0(x)}.$$

On the other hand, by assumption (8.5) and the parabolic Harnack inequality there exists a positive constant C_1 which depends on P_+ , P_0 , y_0 , and y_1 such that

$$(8.11) \quad \begin{aligned} C_1^{-1} k_{P_+}^M(x, y_0, t-1) &\leq k_{P_+}^M(x, y_1, t) \\ &\leq C k_{P_0}^M(x, y_1, t) \leq C C_1 k_{P_0}^M(x, y_0, t+1) \quad \forall x \in M, t > T(x). \end{aligned}$$

Moreover, by (7.8) we have

$$(8.12) \quad \lim_{t \rightarrow \infty} \frac{k_{P_+}^M(x, y_0, t-1)}{k_{P_+}^M(x, y_0, t)} = 1, \quad \text{and} \quad \lim_{t \rightarrow \infty} \frac{k_{P_0}^M(x, y_0, t+1)}{k_{P_0}^M(x, y_0, t)} = 1 \quad \forall x \in M.$$

Therefore, (8.11) and (8.12) imply that there exists $C_0 > 0$ such that

$$(8.13) \quad k_{P_+}^M(x, y_0, t) \leq C_0 k_{P_0}^M(x, y_0, t) \quad \forall x \in M, t > T(x).$$

Consequently, (8.10) and (8.13) imply that

$$u_+(x) \leq C_0 u_0(x) = \tilde{C}_0 \varphi(x) \quad \forall x \in M.$$

Therefore, using (8.4) we obtain

$$(8.14) \quad (u_+)^2(x) m_+(x) A_+(x) \leq C_2 \varphi^2(x) m_0(x) A_0(x) \quad \text{for a.e. } x \in M,$$

where $C_2 > 0$ is a positive constant. Thus, Theorem 2.11 implies that P_+ is critical in M which is a contradiction. The last statement of the theorem follows from the parabolic Harnack inequality and parabolic regularity. \square

By the generalized maximum principle, assumption (8.5) in Theorem 8.4 is satisfied with $C = 1$ if $P_+ = P_0 + V$, where P_0 is a critical operator on M and V is any *nonnegative* potential. Note that if the potential is in addition nontrivial, then P_+ is indeed subcritical in M . Therefore, we have

Corollary 8.5 ([24]). *Let P_0 be a symmetric operator of the form (8.3) which is critical in M , and let $P_+ := P_0 + V_1$, where V_1 is a nonzero nonnegative potential. Then*

$$(8.15) \quad \lim_{t \rightarrow \infty} \frac{k_{P_+}^M(x, y, t)}{k_{P_0}^M(x, y, t)} = 0$$

locally uniformly in $M \times M$.

Next, we discuss the nonsymmetric case. We study two cases where Davies' conjecture implies Conjecture 8.1. First, we show that in the nonsymmetric case, the result of Corollary 8.5 for positive perturbations of a critical operator P_0 still holds provided that the validity of Davies' conjecture (Conjecture 7.1) is assumed instead of the symmetry hypothesis. More precisely, we have

Theorem 8.6 ([24]). *Let P_0 be a critical operator in M , and let $P_+ = P_0 + V$, where V is any nonzero nonnegative potential on M . Assume*

that Davies' conjecture (Conjecture 7.1) holds true for both $k_{P_0}^M$ and $k_{P_+}^M$. Then

$$(8.16) \quad \lim_{t \rightarrow \infty} \frac{k_{P_+}^M(x, y, t)}{k_{P_0}^M(x, y, t)} = 0$$

locally uniformly in $M \times M$.

Proof. Recall that in light of Remark 8.2 we assume that $\lambda_0(P_+, M) = 0$. Suppose to the contrary that for some $x_0, y_0 \in M$ there exists a sequence $\{t_n\}$ such that $t_n \rightarrow \infty$ and

$$(8.17) \quad \lim_{n \rightarrow \infty} \frac{k_{P_+}^M(x_0, y_0, t_n)}{k_{P_0}^M(x_0, y_0, t_n)} = K > 0.$$

Consider the functions v_+ and v_0 defined by

$$v_+(x, t) := \frac{k_{P_+}^M(x, y_0, t)}{k_{P_+}^M(x_0, y_0, t)}, \quad v_0(x, t) := \frac{k_{P_0}^M(x, y_0, t)}{k_{P_0}^M(x_0, y_0, t)} \quad x \in M, t > 0.$$

By our assumption,

$$\lim_{t \rightarrow \infty} v_+(x, t) = u_+(x), \quad \lim_{t \rightarrow \infty} v_0(x, t) = u_0(x),$$

where $u_+ \in \mathcal{C}_{P_+}(M)$ and $u_0 \in \mathcal{C}_{P_0}(M)$.

On the other hand, by the generalized maximum principle

$$(8.18) \quad \frac{k_{P_+}^M(x, y_0, t)}{k_{P_0}^M(x, y_0, t)} \leq 1.$$

Therefore,

$$(8.19) \quad \frac{k_{P_+}^M(x_0, y_0, t_n)}{k_{P_0}^M(x_0, y_0, t_n)} \times \frac{\frac{k_{P_+}^M(x, y_0, t_n)}{k_{P_+}^M(x_0, y_0, t_n)}}{\frac{k_{P_0}^M(x, y_0, t_n)}{k_{P_0}^M(x_0, y_0, t_n)}} = \frac{k_{P_+}^M(x, y_0, t_n)}{k_{P_0}^M(x, y_0, t_n)} \leq 1.$$

Letting $n \rightarrow \infty$ we obtain

$$(8.20) \quad Ku_+(x) \leq u_0(x) \quad x \in M.$$

It follows that $v(x) := u_0(x) - Ku_+(x)$ is a nonnegative supersolution of the equation $P_0 u = 0$ in M which is not a solution. In particular, $v \neq 0$. By the strong maximum principle $v(x)$ is a strictly positive supersolution of the equation $P_0 u = 0$ in M which is not a solution. This contradicts the criticality of P_0 in M . \square

The second nonsymmetric result concerns semismall perturbations.

Theorem 8.7 ([24]). *Let P_+ and $P_0 = P_+ + V$ be a subcritical operator and a critical operator in M , respectively. Suppose that V is a semismall perturbation of the operator P_+ in M . Assume further that Davies' conjecture (Conjecture 7.1) holds true for both $k_{P_0}^M$ and $k_{P_+}^M$ and that (8.5) holds true. Then*

$$(8.21) \quad \lim_{t \rightarrow \infty} \frac{k_{P_+}^M(x, y, t)}{k_{P_0}^M(x, y, t)} = 0$$

locally uniformly in $M \times M$.

Proof. Recall that we (may) assume that $\lambda_0(P_+, M) = 0$. Assume to the contrary that for some $x_0, y_0 \in M$ there exists a sequence $\{t_n\}$ such that $t_n \rightarrow \infty$ and

$$(8.22) \quad \lim_{n \rightarrow \infty} \frac{k_{P_+}^M(x_0, y_0, t_n)}{k_{P_0}^M(x_0, y_0, t_n)} = K > 0.$$

Consider the functions v_+ and v_0 defined by

$$(8.23) \quad v_+(x, t) := \frac{k_{P_+}^M(x, y_0, t)}{k_{P_+}^M(x_0, y_0, t)}, \quad v_0(x, t) := \frac{k_{P_0}^M(x, y_0, t)}{k_{P_0}^M(x_0, y_0, t)} \quad x \in M, t > 0.$$

By our assumption,

$$\lim_{t \rightarrow \infty} v_+(x, t) = u_+(x), \quad \lim_{t \rightarrow \infty} v_0(x, t) = u_0(x),$$

where $u_+ \in \mathcal{C}_{P_+}(M)$ and $u_0 \in \mathcal{C}_{P_0}(M)$.

On the other hand, by assumption (8.5) we have for $t > T(x)$

$$(8.24) \quad \frac{k_{P_+}^M(x, y_0, t)}{k_{P_0}^M(x, y_0, t)} = \frac{k_{P_+}^M(x, y_1, t)}{k_{P_0}^M(x, y_1, t)} \times \frac{\frac{k_{P_+}^M(x, y_0, t)}{k_{P_+}^M(x, y_1, t)}}{\frac{k_{P_0}^M(x, y_0, t)}{k_{P_0}^M(x, y_1, t)}} \leq C \frac{k_{P_+}^M(x, y_0, t)}{k_{P_+}^M(x, y_1, t)} \times \frac{k_{P_0}^M(x, y_1, t)}{k_{P_0}^M(x, y_0, t)}.$$

By our assumption on Davies' conjecture, we have for a fixed x

$$(8.25) \quad \lim_{t \rightarrow \infty} \frac{k_{P_+}^M(x, y_0, t)}{k_{P_+}^M(x, y_1, t)} = \frac{u_+^*(y_0)}{u_+^*(y_1)}, \quad \lim_{t \rightarrow \infty} \frac{k_{P_0}^M(x, y_1, t)}{k_{P_0}^M(x, y_0, t)} = \frac{u_0^*(y_1)}{u_0^*(y_0)},$$

where u_+^* and u_0^* are positive solutions of the equation $P_+^* u = 0$ and $P_0^* u = 0$ in M , respectively. By the elliptic Harnack inequality there exists a positive constant C_1 (depending on P_+^*, P_0^*, y_0, y_1 but not on x) such that

$$(8.26) \quad \frac{u_+^*(y_0)}{u_+^*(y_1)} \leq C_1, \quad \frac{u_0^*(y_1)}{u_0^*(y_0)} \leq C_1.$$

Therefore, (8.24) and (8.26) imply that

$$(8.27) \quad \frac{k_{P_+}^M(x, y_0, t_n)}{k_{P_0}^M(x, y_0, t_n)} \leq 2CC_1^2$$

for n sufficiently large (which might depend on x).

Therefore,

$$(8.28) \quad \frac{k_{P_+}^M(x_0, y_0, t_n)}{k_{P_0}^M(x_0, y_0, t_n)} \times \frac{\frac{k_{P_+}^M(x, y_0, t_n)}{k_{P_+}^M(x_0, y_0, t_n)}}{\frac{k_{P_0}^M(x, y_0, t_n)}{k_{P_0}^M(x_0, y_0, t_n)}} = \frac{k_{P_+}^M(x, y_0, t_n)}{k_{P_0}^M(x, y_0, t_n)} \leq 2CC_1^2.$$

Letting $n \rightarrow \infty$ and using (8.22) and (8.23), we obtain

$$(8.29) \quad Ku_+(x) \leq 2CC_1^2 u_0(x) \quad x \in M.$$

On the other hand, since V is a semismall perturbation of P_+ in M , Theorem 2.13 implies that $u_0(x) \asymp G_{P_+}^M(x, y_0)$ in $M \setminus \overline{B(y_0, \delta)}$, with some positive δ . Consequently,

$$(8.30) \quad u_+(x) \leq C_2 G_{P_+}^M(x, y_0) \quad x \in M \setminus \overline{B(y_0, \delta)}$$

for some $C_2 > 0$. In other words, u_+ is a global positive solution of the equation $P_+ u = 0$ in M which has minimal growth in a neighborhood of infinity in M . Therefore u_+ is a ground state of the equation $P_+ u = 0$ in M , but this contradicts the subcriticality of P_+ in M . \square

9. ON THE EQUIVALENCE OF HEAT KERNELS

In this section we study a general question concerning the equivalence of heat kernels which in turn will give sufficient conditions for the validity of the boundedness condition (8.5) that is assumed in theorems 8.4 and 8.7.

Definition 9.1. Let P_i , $i = 1, 2$, be two elliptic operators of the form (2.1) that are defined on M and satisfy $\lambda_0(P_i, M) = 0$ for $i = 1, 2$. We say that the heat kernels $k_{P_1}^M(x, y, t)$ and $k_{P_2}^M(x, y, t)$ are *equivalent* (respectively, *semiequivalent*) if

$$k_{P_1}^M \asymp k_{P_2}^M \quad \text{on } M \times M \times (0, \infty)$$

(respectively,

$$k_{P_1}^M(\cdot, y_0, \cdot) \asymp k_{P_2}^M(\cdot, y_0, \cdot) \quad \text{on } M \times (0, \infty)$$

for some fixed $y_0 \in M$).

There is an intensive literature dealing with (almost optimal) conditions under which two positive (minimal) Green functions are equivalent or semiequivalent (see [3, 43, 46, 50] and the references therein). On the other hand, sufficient conditions for the equivalence of heat kernels are known only in a few cases (see [39, 40, 67]). In particular, it seems that the answer to the following conjecture is not known.

Conjecture 9.2 ([24]). *Let P_1 and P_2 be two subcritical operators of the form (2.1) that are defined on a Riemannian manifold M such that $P_1 = P_2$ outside a compact set in M , and $\lambda_0(P_i, M) = 0$ for $i = 1, 2$. Then $k_{P_1}^M$ and $k_{P_2}^M$ are equivalent.*

Remark 9.3. Suppose that P_1 and P_2 satisfy the assumption of Conjecture 9.2. B. Devyver and the author proved recently that there exists $C > 0$ such that

$$(9.1) \quad C^{-1}G_{P_2-\lambda}^M(x, y) \leq G_{P_1-\lambda}^M(x, y) \leq CG_{P_2-\lambda}^M(x, y) \quad \forall x, y \in M, \text{ and } \lambda \leq 0.$$

Clearly, by (1.7), the above estimate (9.1) is a necessary condition for the validity of Conjecture 9.2.

It is well known that certain 3- G inequalities imply the equivalence of Green functions, and the notions of small and semismall perturbations is based on this fact. Moreover, small (respectively, semismall) perturbations are sufficient conditions and in some sense also necessary conditions for the equivalence (respectively, semiequivalence) of the Green functions [43, 46, 50]. Therefore, it is natural to introduce an analog definition for heat kernels (cf. [67]).

Definition 9.4. Let P be a subcritical operator in M . We say that a potential V is a k -bounded perturbation (respectively, k -semibounded perturbation) with respect to the heat kernel $k_P^M(x, y, t)$ if there exists a positive constant C such that the following 3- k inequality is satisfied:

$$(9.2) \quad \int_0^t \int_M k_P^M(x, z, t-s) |V(z)| k_P^M(z, y, s) \, dz \, ds \leq C k_P^M(x, y, t)$$

for all $x, y \in M$ (respectively, for a fixed $y \in M$, and all $x \in M$) and $t > 0$.

The following result shows that the notion of k -(semi)boundedness is naturally related to the (semi)equivalence of heat kernels.

Theorem 9.5 ([24]). *Let P be a subcritical operator in M , and assume that the potential V is k -bounded perturbation (respectively, k -semibounded perturbation) with respect to the heat kernel $k_P^M(x, y, t)$. Then there exists $c > 0$ such that for any $|\varepsilon| < c$ the heat kernels*

$k_{P+\varepsilon V}^M(x, y, t)$ and $k_P^M(x, y, t)$ are equivalent (respectively, semiequivalent).

Proof. Consider the iterated kernels

$$k_P^{(j)}(x, y, t) := \begin{cases} k_P^M(x, y, t) & j = 0, \\ \int_0^t \int_M k_P^{(j-1)}(x, z, t-s) V(z) k_P^M(z, y, s) dz ds & j \geq 1. \end{cases}$$

Using (9.2) and an induction argument, it follows that

$$\begin{aligned} \sum_{j=0}^{\infty} |\varepsilon|^j |k_P^{(j)}(x, y, t)| \\ \leq (1 + C|\varepsilon| + C^2|\varepsilon|^2 + \dots) k_P^M(x, y, t) = \frac{1}{1 - C|\varepsilon|} k_P^M(x, y, t) \end{aligned}$$

provided that $|\varepsilon| < C^{-1}$. Consequently, for such ε the Neumann series

$$\sum_{j=0}^{\infty} (-\varepsilon)^j k_P^{(j)}(x, y, t)$$

converges locally uniformly in $M \times M \times \mathbb{R}_+$ to $k_{P+\varepsilon V}^M(x, y, t)$ which in turn implies that Duhamel's formula

$$(9.3) \quad k_{P+\varepsilon V}^M(x, y, t) = k_P^M(x, y, t) - \varepsilon \int_0^t \int_M k_P^M(x, z, t-s) V(z) k_{P+\varepsilon V}^M(z, y, s) dz ds$$

is valid. Moreover, we have

$$k_{P+\varepsilon V}^M(x, y, t) \leq \frac{1}{1 - C|\varepsilon|} k_P^M(x, y, t).$$

The lower bound

$$C_1 k_P^M(x, y, t) \leq k_{P+\varepsilon V}^M(x, y, t)$$

(for $|\varepsilon|$ small enough) follows from the upper bound using (9.3) and (9.2). \square

Corollary 9.6 ([24]). *Assume that P and V satisfy the conditions of Theorem 9.5, and suppose further that V is nonnegative. Then there exists $c > 0$ such that for any $\varepsilon > -c$ the heat kernels $k_{P+\varepsilon V}^M(x, y, t)$ and $k_P^M(x, y, t)$ are equivalent (respectively, semiequivalent).*

Proof. By Theorem 9.5 the heat kernels $k_{P+\varepsilon V}^M(x, y, t)$ and $k_P^M(x, y, t)$ are equivalent (respectively, semiequivalent) for any $|\varepsilon| < c$.

Recall that by the generalized maximum principle,

$$(9.4) \quad k_{P+\varepsilon V}^M(x, y, t) \leq k_P^M(x, y, t) \quad \forall \varepsilon > 0.$$

On the other hand, given a potential W (not necessarily of definite sign), and $0 \leq \alpha \leq 1$, denote $P_\alpha := P + \alpha W$, and assume that $P_j \geq 0$ in M for $j = 0, 1$. Then for $0 \leq \alpha \leq 1$ we have $P_\alpha \geq 0$ in M , and the following inequality holds [47]

$$(9.5) \quad k_{P_\alpha}^M(x, y, t) \leq [k_{P_0}^M(x, y, t)]^{1-\alpha} [k_{P_1}^M(x, y, t)]^\alpha \quad \forall x, y \in M, t > 0.$$

Using (9.4) and (9.5) we obtain the desired equivalence $k_{P+\varepsilon V}^M \asymp k_P^M$ also for $\varepsilon \geq c$. \square

Finally we have:

Theorem 9.7 ([24]). *Let P_0 be a critical operator in M . Assume that $V = V_+ - V_-$ is a potential such that $V_\pm \geq 0$ and $P_+ := P_0 + V$ is subcritical in M .*

Assume further that V_- is k -semibounded perturbation with respect to the heat kernel $k_{P_+}^M(x, y_1, t)$. Then condition (8.5) is satisfied uniformly in x . That is, there exist positive constants C and T such that

$$(9.6) \quad k_{P_+}^M(x, y_1, t) \leq C k_{P_0}^M(x, y_1, t) \quad \forall x \in M, t > T.$$

Proof. It follows from Corollary 9.6, that the heat kernels $k_{P_+}^M(x, y_1, t)$ and $k_{P_++V_-}^M(x, y_1, t)$ are semiequivalent. Note that $P_+ + V_- = P_0 + V_+$. Therefore we have

$$\begin{aligned} C^{-1} k_{P_+}^M(x, y_1, t) &\leq k_{P_++V_-}^M(x, y_1, t) \\ &= k_{P_0+V_+}^M(x, y_1, t) \leq k_{P_0}^M(x, y_1, t) \quad \forall x \in M, t > 0. \end{aligned}$$

\square

Acknowledgments

The author acknowledges the support of the Israel Science Foundation (grant 963/11) founded by the Israeli Academy of Sciences and Humanities.

REFERENCES

- [1] S. Agmon, On positivity and decay of solutions of second order elliptic equations on Riemannian manifolds, in "Methods of Functional Analysis and Theory of Elliptic Equations", pp. 19–52, (ed. D. Greco), Liguori, Naples, 1983.
- [2] ———, *private communication*.
- [3] A. Ancona, First eigenvalues and comparison of Green's functions for elliptic operators on manifolds or domains, *J. Anal. Math.* **72** (1997), 45–92.

- [4] J-P Anker, P. Bougerol, and T. Jeulin, The infinite Brownian loop on symmetric space, *Rev. Mat. Iberoamericana* **18** (2002), 41–97.
- [5] W. Arendt, and C. J. K. Batty, Exponential stability of a diffusion equation with absorption, *Differential Integral Equations* **6** (1993), 1009–1024.
- [6] D. G. Aronson, Non-negative solutions of linear parabolic equations, *Ann. Scuola Norm. Sup. Pisa* (3) **22** (1968), 607–694.
- [7] R. Bañuelos, and B. Davis, Heat kernel, eigenfunctions, and conditioned Brownian motion in planar domains, *J. Funct. Anal.* **84** (1989), 188–200.
- [8] N. H. Bingham, C. M. Goldie, and J. L. Teugels, “Regular Variation”, *Encyclopedia of Mathematics and its Applications* **27**, Cambridge University Press, Cambridge, 1989.
- [9] K. Burdzy and T. S. Salisbury, On minimal parabolic functions and time-homogeneous parabolic h -transforms, *Trans. Amer. Math. Soc.* **351** (1999), 3499–3531.
- [10] O. Calin, D. C. Chang, K. Furutani, and C. Iwasaki, “Heat kernels for elliptic and sub-elliptic operators, Methods and techniques”, *Applied and Numerical Harmonic Analysis*. Birkhäuser/Springer, New York, 2011.
- [11] I. Chavel, “Isoperimetric Inequalities. Differential Geometric and Analytic Perspectives”, *Cambridge Tracts in Mathematics* **145**, Cambridge University Press, Cambridge, 2001.
- [12] I. Chavel and L. Karp, Large time behavior of the heat kernel: the parabolic λ -potential alternative, *Comment. Math. Helv.* **66** (1991), 541–556.
- [13] K. L. Chung, “Markov Chains with Stationary Transition Probabilities” 2nd ed., *Die Grundlehren der mathematischen Wissenschaften*, **104**, Springer-Verlag, New York, 1967.
- [14] P. Collet, S. Martínez, and J. San Martín, Ratio limit theorems for a Brownian motion killed at the boundary of a Benedicks domain, *Ann. Probab.*, **27** (1999), 1160–1182.
- [15] ———, Asymptotic of the heat kernel in general Benedicks domains, *Probab. Theory Related Fields* **125** (2003), 350–364.
- [16] C. Constantinescu, and A. Cornea, “Potential Theory on Harmonic Spaces”, *Die Grundlehren der Mathematischen Wissenschaften* **158**, Springer-Verlag, New York-Heidelberg, 1972.
- [17] T. Coulhon, Heat kernels on non-compact Riemannian manifolds: a partial survey, in “Séminaire de Théorie Spectrale et Géométrie”, pp. 167–187, *Sémin. Théor. Spectr. Géom.* **15**, Univ. Grenoble I, Saint-Martin-d’Hères, 1997.
- [18] ———, Large time behaviour of heat kernels on Riemannian manifolds: fast and slow decays, *Journées “Équations aux Dérivées Partielles”* (Saint-Jean-de-Monts, 1998), Exp. No. II, 12 pp., Univ. Nantes, Nantes, 1998.
- [19] T. Coulhon, and A. Grigor’yan, On-diagonal lower bounds for heat kernels on non-compact manifolds and Markov chains, *Duke Math. J.* **89** (1997), 133–199.
- [20] E. B. Davies, “Heat Kernels and Spectral Theory”, *Cambridge Tracts in Mathematics* **92**, Cambridge University Press, Cambridge, 1990.
- [21] ———, Heat kernel bounds, conservation of probability and the Feller property, *J. Anal. Math.* **58** (1992), 99–119.
- [22] ———, Non-Gaussian aspects of heat kernel behaviour, *J. London Math. Soc.* (2) **55** (1997), 105–125.

- [23] J. L. Doob, “Classical potential theory and its probabilistic counterpart”, Grundlehren der Mathematischen Wissenschaften **262**, Springer-Verlag, New York, 1984.
- [24] M. Fraas, D. Krejčířík, and Y. Pinchover, On some strong ratio limit theorems for heat kernels, *Disc. Contin. Dyn. Syst.* **28** (2010), 495–509.
- [25] A. Grigor’yan, Estimates of heat kernels on Riemannian manifolds, in “Spectral Theory and Geometry”, pp. 140–225, London Math. Soc. Lecture Note Ser. **273**, Cambridge Univ. Press, Cambridge, 1999.
- [26] ———, Analytic and geometric background of recurrence and non-explosion of the Brownian motion on Riemannian manifolds, *Bull. Amer. Math. Soc.* **36** (1999), 135–249.
- [27] ———, “Heat Kernel and Analysis on Manifolds”, AMS/IP Studies in Advanced Mathematics, 47, American Mathematical Society, Providence, RI, International Press, Boston, MA, 2009.
- [28] A. Grigor’yan, and L. Saloff-Coste, Heat kernel on manifolds with ends, *Ann. Inst. Fourier (Grenoble)* **59** (2009), 1917–1997.
- [29] R. Z. Has’minskii, “Stochastic Stability of Differential Equations”, Monographs and Textbooks on Mechanics of Solids and Fluids: Mechanics and Analysis, **7**, Sijthoff & Noordhoff, Alphen aan den Rijn, 1980.
- [30] K. Ishige, and M. Murata, Uniqueness of nonnegative solutions of the Cauchy problem for parabolic equations on manifolds or domains, *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)* **30** (2001), 171–223.
- [31] P. Li, Large time behavior of the heat equation on complete manifolds with nonnegative Ricci curvature, *Ann. of Math. (2)* **124** (1986), 1–21.
- [32] M. Keller, D. Lenz, H. Vogt, and R. Wojciechowski, Note on basic features of large time behaviour of heat kernels. arXiv: 1101.0373.
- [33] J. F. C. Kingman, Ergodic properties of continuous-time Markov processes and their discrete skeletons, *Proc. London Math. Soc. (3)* **13** (1963), 593–604.
- [34] W. Kirsch, and B. Simon, Approach to equilibrium for a forced Burgers equation, *J. Evol. Equ.* **1** (2001), 411–419.
- [35] G. Kozma, A graph counterexample to Davies’ conjecture. arXiv:1111.4593.
- [36] D. Krejčířík, and E. Zuazua, The Hardy inequality and the heat equation in twisted tubes, *J. Math. Pures Appl. (9)* **94** (2010), 277–303.
- [37] P. Li, “Geometric Analysis” (Cambridge Studies in Advanced Mathematics; 134), Cambridge University Press, Cambridge, 2012.
- [38] V. Lin and Y. Pinchover, Manifolds with group actions and elliptic operators, *Memoirs Amer. Math. Soc.* **112**, (1994), 1–78.
- [39] V. Liskevich, and Y. Semenov, Two-sided estimates of the heat kernel of the Schrödinger operator, *Bull. London Math. Soc.*, **30** (1998), 596–602.
- [40] P. D. Milman, and Yu. A. Semenov, Heat kernel bounds and desingularizing weights, *J. Funct. Anal.*, **202** (2003), 1–24.
- [41] M. Murata, Positive solutions and large time behaviors of Schrödinger semi-groups, Simon’s problem, *J. Funct. Anal.*, **56** (1984), 300–310.
- [42] ———, Structure of positive solutions to $(-\Delta + V)u = 0$ in \mathbb{R}^n , *Duke Math. J.* **53** (1986), 869–943.
- [43] ———, Semismall perturbations in the Martin theory for elliptic equations, *Israel J. Math.*, **102** (1997), 29–60.

- [44] ———, Martin boundaries of elliptic skew products, semismall perturbations, and fundamental solutions of parabolic equations, *J. Funct. Anal.* **194** (2002), 53–141.
- [45] ———, Integral representations of nonnegative solutions for parabolic equations and elliptic Martin boundaries, *J. Funct. Anal.* **245** (2007), 177–212.
- [46] Y. Pinchover, Criticality and ground states for second-order elliptic equations, *J. Differential Equations*, **80** (1989), 237–250.
- [47] ———, On criticality and ground states of second order elliptic equations. II, *J. Differential Equations* **87** (1990), 353–364.
- [48] ———, Large time behavior of the heat kernel and the behavior of the Green function near criticality for nonsymmetric elliptic operators, *J. Funct. Anal.* **104** (1992), 54–70.
- [49] ———, On nonexistence of any λ_0 -invariant positive harmonic function, a counterexample to Stroock’s conjecture, *Comm. Partial Differential Equations* **20** (1995), 1831–1846.
- [50] ———, Maximum and anti-maximum principles and eigenfunctions estimates via perturbation theory of positive solutions of elliptic equations, *Math. Ann.*, **314** (1999), 555–590.
- [51] ———, Large time behavior of the heat kernel, *J. Funct. Anal.* **206** (2004), 191–209.
- [52] ———, On Davies’ conjecture and strong ratio limit properties for the heat kernel, in “Potential Theory in Matsue”, Proceedings of the International Workshop on Potential Theory, 2004, pp.339–352, (eds. H. Aikawa, et al.), Advanced Studies in Pure Mathematics **44**, Mathematical Society of Japan, Tokyo, 2006.
- [53] ———, A Liouville-type theorem for Schrödinger operators, *Comm. Math. Phys.*, **272** (2007), 75–84.
- [54] R. G. Pinsky, “Positive Harmonic Function and Diffusion”, Cambridge University Press, Cambridge, 1995.
- [55] F. O. Porper, and S. D. Eidel’man, Two-sided estimates of the fundamental solutions of second-order parabolic equations and some applications of them, *Uspekhi Mat. Nauk* **39** (1984), 107–156.
- [56] V. C. Repnikov, S. D. Eidel’man, Necessary and sufficient conditions for establishing a solution to the Cauchy problem, *Soviet Math. Dokl.* **7** (1966), 388–391.
- [57] D. W. Robinson, “Elliptic Operators and Lie Groups”, Oxford Mathematical Monographs, Oxford University Press, New York, 1991.
- [58] B. Simon, “Functional integration and quantum physics”, Pure and Applied Mathematics, 86, Academic Press, Inc., New York-London, 1979.
- [59] ———, Schrödinger semigroups, *Bull. Amer. Math. Soc.* **7** (1982), 447–526.
- [60] ———, Large time behavior of the heat kernel: on a theorem of Chavel and Karp, *Proc. Amer. Math. Soc.* **118** (1993), 513–514.
- [61] U. Stadtmüller, and R. Trautner, Ratio Tauberian theorems for Laplace transforms without monotonicity assumptions, *Quart. J. Math. Oxford Ser. (2)* **36** (1985), 363–381.
- [62] S. R. S. Varadhan, “Lectures on Diffusion Problems and Partial Differential Equations”, Tata Institute of Fundamental Research **64**, Springer-Verlag, Berlin, 1980.

- [63] N. Th. Varopoulos, L. Saloff-Coste, and T. Coulhon, “Analysis and Geometry on Groups”, Cambridge Tracts in Mathematics **100**, Cambridge University Press, Cambridge, 1992.
- [64] W. Woess, “Random Walks on Infinite Graphs and Groups”, Cambridge Tracts in Mathematics **138**, Cambridge University Press, Cambridge, 2000.
- [65] K. Wong, Large time behavior of Dirichlet heat kernels on unbounded domains above the graph of a bounded Lipschitz function, *Glas. Mat. Ser. III* **41 (61)** (2006), 177–186.
- [66] Q. S. Zhang, Large time behavior of Schrödinger heat kernels and applications, *Comm. Math. Phys.* **210** (2000), 371–398.
- [67] ———, A sharp comparison result concerning Schrödinger heat kernels, *Bull. London Math. Soc.*, **35** (2003), 461–472.

DEPARTMENT OF MATHEMATICS, TECHNION - ISRAEL INSTITUTE OF TECHNOLOGY, HAIFA 32000, ISRAEL

E-mail address: `pincho@techunix.technion.ac.il`